

# Radiocarbon dates from abandoned penguin colonies in the Antarctic Peninsula region

STEVEN D. EMSLIE

Department of Biological Sciences, University of North Carolina, Wilmington, NC 28403, USA  
emslies@uncwil.edu

**Abstract:** Sixty-three radiocarbon dates on organic remains from 21 abandoned colonies of chinstrap (*Pygoscelis antarctica*) and Adélie (*P. adeliae*) penguins on 12 islands in the Antarctic Peninsula region are evaluated for determining the occupation history of penguins in this region. This record also provides a means for assessing sea-level change, glacial advances and retreats, and population responses by penguins to these events. All conventional dates were corrected for the marine-carbon reservoir effect by applying a  $\Delta R = 700 \pm 50$  BP and marine calibration curves. The 63 calibrated dates give  $2\sigma$  ranges (95% confidence intervals) from modern to 5990 yr BP. These dates indicate progressively older occupations from north to south along the Antarctic Peninsula. No sites older than approximately 540 BP occur in the northern peninsula, either because they have not yet been found or older sites have been destroyed by solifluction and glacial scouring. Three dates from one locality near Rothera Point, Adelaide Island, also were calibrated with a  $\Delta R = 750 \pm 50$  and  $800 \pm 50$  BP. No difference was found between calibrated dates using these two other  $\Delta R$  values, indicating that local corrections for variation in upwelling intensity may not be necessary.

Received 28 November 2000, accepted 22 May 2001

**Key words:** abandoned colonies, Adélie penguin, chinstrap penguin, occupation history, radiocarbon dates

## Introduction

Abandoned penguin colonies in Antarctica provide a source of well-preserved organic remains that date from hundreds to thousands of years old. These remains not only include bones, tissue, and eggshell fragments of penguins that once occupied these sites, but also hard parts from their prey that accumulated in the sediments from penguin guano. The high concentration of these organic remains in ornithogenic soils allows study of the occupation history and diet of penguins in the past. A radiocarbon chronology of the abandoned sites also can be compared to the palaeoclimatic record in Antarctica, based on ice cores and marine sediments, to investigate ecological responses by penguins to episodes of climate change (Smith *et al.* 1999).

Early research on abandoned colonies concentrated on nutrient cycling and chemistry of the ornithogenic soils that characterize these deposits (Ugolini 1972, Speir & Heine 1982, Speir & Cowling 1984, Speir & Ross 1984, Myrcha *et al.* 1985, Tatur 1987, 1989, Heine & Speir 1989, Tatur & Myrcha 1989, Tatur *et al.* 1997). These soils were first identified and described in the Antarctic by Syroechkovsky (1959) and Tedrow & Ugolini (1966) (see also discussion by Campbell & Claridge 1987). Other studies on abandoned colonies have concentrated on the occupation history of penguins and palaeoclimatic inferences as obtained from radiocarbon dates of bones from ornithogenic soils (Harrington & McKellar 1958, Spellerberg 1970, Stonehouse 1970, Bochenski 1985, Moors *et al.* 1988, Baroni & Orombelli 1991, 1994, Smith *et al.* 1999), or on biochemical analyses of

ancient guano in lake sediments (Zale 1994a, Sun *et al.* 2000). Palaeodietary studies and taphonomy of bone deposits in abandoned colonies of chinstrap (*Pygoscelis antarctica* Forster) and Adélie (*P. adeliae* Hombron & Jacquinot) penguins were initiated by Emslie (1995) and Emslie *et al.* (1998) in the Antarctic Peninsula region.

Accurate interpretation of data from abandoned colonies is dependent on the radiocarbon chronology obtained from organic remains recovered from these sites. Numerous methods have been proposed for correction and calibration of radiocarbon dates of marine organisms in Antarctica, necessary due to the marine-carbon reservoir effect (Harkness 1979, Björck *et al.* 1991, Gordon & Harkness 1992, Emslie 1995, Berkman & Forman 1996, Berkman *et al.* 1998). These methods adjust the conventional radiocarbon age by either subtracting a pre-determined number of years from all dates, or by applying a correction and calibration to each date using a  $\Delta R$  value that incorporates changing differences in reservoir effects between the atmosphere and ocean through time (Stuiver *et al.* 1986, 1998). The former method provides a corrected date in radiocarbon years, while the latter produces a date calibrated to calendar years BP. Both methods are based on radiocarbon dates obtained on modern (pre-1950) specimens of marine organisms that yield ages from 100 s to > 1000 years due to ingestion of old carbon during life. Here, I examine 60 radiocarbon dates on pygoscelid penguin bone, tissue, and prey remains, and three on ornithogenic sediments, from abandoned colonies in the Antarctic Peninsula region. These dates are assessed for their accuracy in estimating the

**Table 1.** Radiocarbon dates (in years BP) on penguin (*Pygoscelis* sp.) bone collagen, eggshell carbonates, or dried dermal tissue and squid (*Psychroteuthis glacialis* Thiele) beaks from test pits (TP) in abandoned colonies or modern (pre-1950) sites. Conventional dates were corrected and calibrated for the marine-carbon reservoir effect using a  $\Delta R = 700 \pm 50$  years (see Emslie 1995) and the MARINE98 database of Stuiver & Reimer (1993). The conventional date, mean corrected date (both in radiocarbon years BP), and calibrated  $2\sigma$  range(s) (95% confidence interval in calendar years BP) are provided for 60 samples from 21 sites on 11 islands. Three dates on modern specimens also are provided. Multiple  $2\sigma$  ranges are provided for corrected dates that intersected two or more regions of the calibration curve. All dates are from Emslie (1995) and Emslie *et al.* (1998) except for UA 1033 (C. Hjort, personal communication 1997), Lu-3101 from Björck *et al.* (1991), and SRR-3784, SRR-3785, and SRR-3786 (R.I. Lewis Smith, personal communication 2001). Dates from the Rothera area and Rothera Point are provided here for the first time. Levels are 5–10 cm deep beginning with level 1 at the surface (0–5 cm) of each test pit (TP) excavated at each site.

Site/sample	Species	Material	Laboratory number	Conventional date	Corrected date	Calibrated range ( $2\sigma$ )
<b>King George Island</b>						
Uchatka, TP 1, level 1	<i>Pygoscelis</i> cf. <i>P. antarctica</i>	bone	CAMS 15626	1300 ± 70	600 ± 85	425–40 or 35–0
	<i>P. antarctica</i>	bone	CAMS 15629	1280 ± 60	580 ± 80	405–40 or 35–0
level 2	<i>P. antarctica</i>	bone	CAMS 15625	1380 ± 60	680 ± 80	475–135
	<i>Pygoscelis</i> cf. <i>P. antarctica</i>	bone	CAMS 15628	1350 ± 50	650 ± 70	445–130
level 3	cf. <i>Pygoscelis</i> sp.	bone	CAMS 15644	1340 ± 60	640 ± 80	450–80
	<i>Pygoscelis</i> sp.	bone	CAMS 14609	1330 ± 70	630 ± 85	450–55 or 15–0
Blue Dyke, TP 1, subsurface	<i>P. antarctica</i>	bone	CAMS 15624	1330 ± 60	630 ± 80	445–65 or 5–0
	<i>P. antarctica</i>	bone	CAMS 14821	1500 ± 60	800 ± 80	540–280
Low Head, surface	<i>P. antarctica</i>	bone	CAMS 15627	1410 ± 60	710 ± 80	495–235
	<i>P. antarctica</i>	bone	CAMS 14817	1440 ± 60	740 ± 80	510–255
<b>Penguin Island</b>						
surface	<i>Pygoscelis</i> cf. <i>P. adeliae</i>	bone	CAMS 14815	1420 ± 60	720 ± 80	500–240
<b>Seymour Island</b>						
surface	<i>Pygoscelis</i> cf. <i>P. adeliae</i>	bone	CAMS 14814	1460 ± 60	760 ± 80	520–265
	<i>Pygoscelis</i> cf. <i>P. adeliae</i>	bone	CAMS 15623	1490 ± 60	790 ± 80	535–275
<b>Devil Island</b>						
surface	<i>P. adeliae</i>	bone	UA 1033	1425 ± 80	725 ± 95	515–145
TP 1, level 3	<i>P. adeliae</i>	bone	Beta 143249	1480 ± 40	780 ± 65	515–285
<b>Litchfield Island</b>						
site 1, TP 2, level 1	<i>P. adeliae</i>	bone	CAMS 42001	1380 ± 50	680 ± 70	465–225, 220–195, or 165–150
	<i>P. adeliae</i>	bone	CAMS 42380	1300 ± 50	600 ± 70	405–55 or 15–0
site 1, TP 2, level 2	<i>P. adeliae</i>	bone	CAMS 42002	1330 ± 50	630 ± 70	425–105
site 2, TP 1, level 1	<i>P. adeliae</i>	bone	CAMS 41997	1230 ± 60	530 ± 80	310–40 or 35–0
site 2, TP 1, level 2	<i>P. adeliae</i>	bone	CAMS 41998	1540 ± 40	840 ± 65	545–325
site 2, TP 2, level 1	<i>P. adeliae</i>	bone	CAMS 41999	1360 ± 50	660 ± 70	450–135
site 2, TP 2, level 2	<i>P. adeliae</i>	bone	CAMS 42000	1240 ± 50	540 ± 70	305–40 or 35–0
modern TP 1, level 3	<i>P. adeliae</i>	bone	CAMS 42011	1170 ± 60	470 ± 80	275–40 or 35–0
<b>Humble Island</b>						
site 1, level 1	<i>P. adeliae</i>	bone	CAMS 42003	1400 ± 50	700 ± 70	475–240
	<i>P. adeliae</i>	bone	CAMS 42004	1360 ± 40	660 ± 65	445–140
	<i>P. adeliae</i>	bone	CAMS 42005	790 ± 50	-	
	<i>P. adeliae</i>	bone	CAMS 42006	1160 ± 50	460 ± 70	265–40 or 35–0
<b>Biscoe Point</b>						
site 1, TP 1, level 1	<i>P. adeliae</i>	bone	CAMS 42007	1540 ± 50	840 ± 70	550–315
	<i>P. adeliae</i>	bone	CAMS 42370	1530 ± 50	830 ± 70	545–310
site 1, TP 1, level 2	<i>P. adeliae</i>	bone	CAMS 42371	1790 ± 40	1090 ± 65	745–540
site 2, level 1	<i>P. adeliae</i>	bone	CAMS 42377	1650 ± 50	950 ± 70	650–460
site 3, level 1	<i>P. adeliae</i>	bone	CAMS 42378	1460 ± 50	760 ± 70	510–270
site 4, level 1	<i>P. adeliae</i>	bone	CAMS 42379	1460 ± 50	760 ± 70	510–270
site 5, TP 1, level 1	<i>P. adeliae</i>	bone	CAMS 42372	1350 ± 50	650 ± 70	445–130
	<i>P. adeliae</i>	bone	CAMS 42373	1180 ± 50	480 ± 70	270–40 or 35–0

occupation history of penguins during the mid to late Holocene in this region. Different methods for correcting and calibrating these dates for the marine reservoir effect also are examined.

## Methods

All organic material collected from the abandoned penguin colonies was obtained through surface collection and

excavation of sites located at Seymour Island (64°15'S, 57°20'W), Devil Island by north Vega Island (63°48'S, 57°15'W), King George Island (62°10'S, 58°30'W), Penguin Island (62°10'S, 58°30'W), Palmer station (64°46'S, 64°04'W) and vicinity at southern Anvers Island, and Rothera station (67°34'S, 68°07'W) and vicinity at south-east Adelaide Island (Fig. 1). Each site was excavated by placing a 1 x 1 m test pit within the pebble concentrations and ornithogenic soils.

**Table 1.** (continued) Radiocarbon dates (in years BP) on penguin (*Pygoscelis* sp.) bone collagen, eggshell carbonates, or dried dermal tissue and squid (*Psychroteuthis glacialis* Thiele) beaks from test pits (TP) in abandoned colonies or modern (pre-1950) sites. Conventional dates were corrected and calibrated for the marine-carbon reservoir effect using a  $\Delta R = 700 \pm 50$  years (see Emslie 1995) and the MARINE98 database of Stuiver & Reimer (1993). The conventional date, mean corrected date (both in radiocarbon years BP), and calibrated  $2\sigma$  range(s) (95% confidence interval in calendar years BP) are provided for 60 samples from 21 sites on 11 islands. Three dates on modern specimens also are provided. Multiple  $2\sigma$  ranges are provided for corrected dates that intersected two or more regions of the calibration curve. All dates are from Emslie (1995) and Emslie *et al.* (1998) except for UA 1033 (C. Hjort, personal communication 1997), Lu-3101 from Björck *et al.* (1991), and SRR-3784, SRR-3785, and SRR-3786 (R.I. Lewis Smith, personal communication 2001). Dates from the Rothera area and Rothera Point are provided here for the first time. Levels are 5–10 cm deep beginning with level 1 at the surface (0–5 cm) of each test pit (TP) excavated at each site.

Site/sample	Species	Material	Laboratory number	Conventional date	Corrected date	Calibrated range ( $2\sigma$ )
<b>Biscoe Point</b>						
site 5, TP 1, level 2	<i>P. adeliae</i>	bone	CAMS 42374	1600 ± 40	900 ± 65	620–425
	<i>P. adeliae</i>	bone	CAMS 42375	1580 ± 50	880 ± 70	615–405
	<i>Psychroteuthis</i>	beak	CAMS 38709	1580 ± 50	880 ± 70	615–405
site 5, TP 1, level 3	<i>P. adeliae</i>	bone	CAMS 42376	1600 ± 50	900 ± 70	625–420
	<i>Psychroteuthis</i>	beak	CAMS 38708	1650 ± 50	950 ± 70	650–460
<b>Cormorant Island</b>						
site 1, TP 1, level 2	<i>Pygoscelis</i> sp.	bone	CAMS 42008	1700 ± 50	1000 ± 70	675–490
	<i>Pygoscelis</i> sp.	bone	CAMS 42009	1760 ± 60	1060 ± 80	745–510
<b>Torgersen Island</b>						
TP1, level 1	<i>P. adeliae</i>	bone	CAMS 42010	1190 ± 40	490 ± 65	270–40 or 35–0
	<i>Psychroteuthis</i>	beak	CAMS 38710	1280 ± 50	580 ± 70	365–40 or 35–0
<b>Ginger Island</b>						
TP 1, level 3	<i>P. adeliae</i>	bone	Beta 141904	1090 ± 40	390 ± 60	-
TP 1, level 4	<i>P. adeliae</i>	bone	Beta 141905	1580 ± 50	880 ± 70	615–405
TP 1, level 5	<i>P. adeliae</i>	bone	Beta 141906	1770 ± 40	1070 ± 60	720–535
TP 1, level 7	<i>P. adeliae</i>	bone	Beta 144171	3130 ± 40	2430 ± 65	2275–1900
<b>Lagoon Island</b>						
site 2, TP 1, level 2	<i>Pygoscelis</i> sp.	bone	Beta 141907	1910 ± 40	1210 ± 60	895–650
site 2, TP 1, level 3	<i>Pygoscelis</i> sp.	bone	Beta 141908	4440 ± 50	3740 ± 70	3835–3470
site 2, TP 1, level 4	<i>Pygoscelis</i> sp.	bone	Beta 141909	4390 ± 50	3690 ± 70	3800–3435
site 2, TP 1, level 5	<i>Pygoscelis</i> sp.	bone	Beta 141910	4550 ± 50	3850 ± 70	3975–3625
site 2, TP 1, level 6	<i>Pygoscelis</i> sp.	bone	Beta 141911	4240 ± 50	3540 ± 70	3585–3265
site 3, TP 1, level 2	<i>Pygoscelis</i> sp.	bone	Beta 141912	2980 ± 50	2280 ± 70	2050–1715
site 3, TP 1, level 3	<i>Pygoscelis</i> sp.	bone	Beta 141913	3510 ± 70	2810 ± 90	2750–2315
site 3, TP 1, level 4	<i>Pygoscelis</i> sp.	bone	Beta 141914	5610 ± 80	4910 ± 95	5460–4950
site 3, TP 1, level 5	<i>Pygoscelis</i> sp.	bone	Beta 141915	5920 ± 50	5220 ± 70	5715–5450
	<i>Pygoscelis</i> sp.	bone	Beta 141916	6080 ± 50	5380 ± 70	5900–5590
site 4, TP 1, level 2	<i>Pygoscelis</i> sp.	bone	Beta 141917	2490 ± 50	1790 ± 70	1490–1220
site 4, TP 1, level 4	<i>Pygoscelis</i> sp.	eggshell	Beta 144170	5550 ± 40	4850 ± 65	5305–4950
north side		sediments	SRR-3786	2680 ± 45	1980 ± 70	1700–1370
<b>Rothera Point, Adelaide Island</b>						
south-west		sediments	SRR-3784	3870 ± 55	3170 ± 70	3160–2770
centre		sediments	SRR-3785	2780 ± 55	2080 ± 70	1820–1500
<b>Modern Specimens</b>						
South Georgia (AMNH 4361)	<i>P. papua</i>	tissue	CAMS 15646	1110 ± 60		
Bay of Whales (AMNH 5041)	<i>P. adeliae</i>	tissue	CAMS 15643	1330 ± 60		
Hope Bay	<i>P. adeliae</i>	bone	Lu-3101	1280 ± 50		

Sediments were removed in 5 or 10 cm levels and washed through three nested screens with mesh sizes of 6.4, 3.2, and 0.025 mm. Organic remains were recovered by sifting through the washed sediments, often with a low-power binocular microscope. Further details on excavation methodology at all sites, except those at Rothera Point, can be found in Emslie (1995) and Emslie *et al.* (1998).

Sites near Rothera Point were excavated in 2000 using

similar methodology. Abandoned colonies were first recognized in this area by R.I. Lewis Smith (British Antarctic Survey 1989). One active colony at Ginger Island (67°45'S, 68°41'W) and three abandoned sites (Sites 2, 3, and 4) at Lagoon Island (67°36'S, 68°14'W) were sampled by excavating 1 m<sup>2</sup> test pits in 5–10 cm levels numbered sequentially with depth. Radiocarbon dates from these sites are reported by excavation level.




Fig. 1. Map of the Antarctic Peninsula region with locations of study sites where abandoned penguin colonies were sampled. Crosses provide latitude and longitude for specific regions.

As noted above, radiocarbon dates of organic remains must be corrected for the marine-carbon reservoir effect and upwelling of old water in the southern ocean (Stuiver *et al.* 1986, Björck *et al.* 1991, Gordon & Harkness 1992, Berkman & Forman 1996, Berkman *et al.* 1998). Rather than correcting each date by subtracting a pre-determined value, Stuiver *et al.* (1986, 1998) provide calibration curves based on the reservoir effects between regional and world oceans through time. Thus, they use a regional offset value ( $\Delta R$ ) that is subtracted from the radiocarbon age. The resulting figure is compared to a calibration curve to estimate the true or corrected age (in calendar years) of the dated material using the Pretoria/Beta Analytic calibration program at Beta Analytic, Inc, Coral Gables, Florida.

Calculation of the  $\Delta R$  depends on radiocarbon dates obtained on modern specimens. These dates will vary by taxa and by geographic location in Antarctica due to differences in species diets, and to variation in the intensity of upwelling and mixing of older waters at continental margins. Moreover, modern samples from which to base a calibration ideally should be from specimens that were collected prior to nuclear testing in the 1950s to avoid additional atmospheric contamination that have influenced carbon ratios in living tissue.

Two specimens of modern, pre-1940 penguins previously were analysed and compared to other published dates by




Fig. 2. Distribution of radiocarbon dates from abandoned and active penguin colonies near Rothera Point, Adelaide Island. Each date is plotted by a pair of points representing the  $2\sigma$  calibrated range (95% confidence interval) after correcting for the marine-carbon reservoir effect. Correlation with glacial history given at the top of the graph is based on data summarized in Ingólfsson *et al.* (1998). Codes and plots refer to: LA4.4 and LA4.2, Lagoon Island Site 4 levels 4 and 2 (plotted with solid diamonds); LA3.5–LA3.2, Lagoon Island Site 3 levels 5–2 (triangles); LA2.6–LA2.2, Lagoon Island Site 2 levels 6–2 (squares); G17, G15, and G14, Ginger Island levels 7, 5, and 4 (circles).

Emslie (1995) to determine a  $\Delta R$  value for pygoscelid penguins in the Antarctic Peninsula region at  $700 \pm 50$  BP. This value was determined after comparing the  $\Delta R$ s of  $715 \pm 50$  for the Bay of Whales, which probably overestimates the value due to more intense upwelling there, vs  $640 \pm 60$  for South Georgia, which underestimates the value as waters are less mixed there than around the Antarctic Peninsula (Emslie 1995). A median value of  $700 \pm 50$  thus appears to be a reasonable estimate and this  $\Delta R$  is applied here to all known dates of penguin bones, tissue, and eggshell from abandoned colonies in the Antarctic Peninsula region.

The method also was applied to squid (*Psychroteuthis glacialis*) beaks identified in the ornithogenic sediments as this species, like pygoscelid penguins, feeds primarily on krill (*Euphausia superba* Dana, Lu & Williams 1994) and therefore may reflect a similar old carbon ratio in its tissues. No modern dates on squid are known from which to assess the accuracy of this calibration. The three dates on sediments from Rothera Point and vicinity were calibrated with this method because the organic material that was dated was largely from penguin guano (R.I. Lewis Smith, personal communication 2001). Zale (1994b) analysed the geochemistry of similar sediments

at Lake Boeckella, Hope Bay, and found that the effect of autochthonous carbon (e.g. from plants) on estimating the true radiocarbon age to be negligible (but see also Björck *et al.* 1991).

It is possible that the  $\Delta R$  applied here for the Antarctic Peninsula region still underestimates the true correction. For example, Emslie (1995) calculates a  $\Delta R$  of  $805 \pm 50$  based on a modern date on Adélie penguin bone from Hope Bay reported by Björck *et al.* (1991). Thus, two other  $\Delta R$  values of  $750 \pm 50$  and  $800 \pm 50$  BP were applied to three conventional dates ranging in age from 1910 to 6080 BP to determine how different  $\Delta R$  values can influence the calibrated results. The two other  $\Delta R$  values were chosen from other published studies on dating marine-based organisms in Antarctica that require an adjustment for the marine-carbon reservoir effect (see Berkman *et al.* 1998).

**Results**

Table I provides a summary of 60 radiocarbon dates from 21 abandoned colonies on 11 islands in the Antarctic Peninsula region, plus three dates on sediments from Rothera Point and vicinity, Adelaide Island. Three modern (pre-1950) dates on penguin bone or tissue also are provided. All specimens were well-preserved and contained amounts of nitrogen and amino acids approaching that of fresh tissue. With few exceptions, multiple radiocarbon dates from one site and test pit are older in progressively deeper levels, or reflect the same age as the level above. This trend is shown by dates both on penguin bones and prey remains (squid beaks).

The  $2\sigma$  calibrated range for each date using a  $\Delta R = 700 \pm 50$  BP indicates that penguin occupation of the King George, Penguin, Devil and Seymour Island sites in the northern Antarctic Peninsula region extends from modern to 540 BP (Table I). Differences in chinstrap vs Adélie penguin occupation of this region were discussed by Emslie (1995). The  $2\sigma$  calibrated range for sites in the Palmer station region is slightly longer, from modern to 745 BP, with all sites representing only Adélie penguin occupation until the recent expansion of chinstrap and gentoo (*Pygoscelis papua* Forster) penguins into the region (Emslie *et al.* 1998, Smith *et al.* 1999). At Rothera Station, the  $2\sigma$  calibrated range is more extensive, from modern to 5900 BP, with all occupations presumably representing Adélie penguins (Table I).

The 21 dated localities form a rough transect from north to south along the Antarctic Peninsula (Fig. 1) and the calibrated dates indicate progressively older occupation southwards along this transect, with the oldest dates from Lagoon Island near Rothera Point. Some of these dates also indicate multiple occupation periods of the same colonies (sites 2–4, Fig. 2). General palaeoclimatic trends for the Antarctic Peninsula (summarized by Ingólfsson *et al.* 1998) correspond with the occupation history of penguins in the Rothera region (Fig. 2).

Three dates (Beta 141907, 141911, and 141916) calibrated with multiple  $\Delta R$  values ( $= 700 \pm 50$ ,  $750 \pm 50$ , and  $800 \pm 50$

BP) indicate broad overlap and no differences in  $2\sigma$  calibrated ranges (Table II).

**Discussion**

Berkman *et al.* (1998) summarized all published methods for dating marine organisms in Antarctica except for that of Emslie (1995), who provided two additional dates on pre-1950 penguin tissue or bone from sub-Antarctic South Georgia and the Bay of Whales, Ross Sea. These dates were used to estimate a  $\Delta R = 700 \pm 50$  for the Antarctic Peninsula region (Emslie 1995). However, Baroni & Orombelli (1991) used a  $\Delta R = 779 \pm 60$  to correct dates from Victoria Land, Ross Sea, while another modern date from Hope Bay in the Peninsula indicated a  $\Delta R = 805 \pm 50$  should be used in that region. Thus, a  $\Delta R$  correction may be subject to local variables including the intensity of upwelling of water containing old carbon (Emslie 1995). Data presented here, however, indicate that application of three  $\Delta R$ s (700, 750, and  $800 \pm 50$  BP) on radiocarbon dates from the same region provides overlapping results and few differences in interpreting the age of the sites (Table II). These results suggest that local variations in upwelling may make little difference in interpreting the true age of pygoscelid penguin remains in Antarctica.

When comparing the data from Rothera with the glacial history record, the estimated calibrated dates seem to match the palaeoclimatic record (Fig. 2). This record, summarized by Ingólfsson *et al.* (1998), indicates that a warming trend and episode of deglaciation occurred at about 6000 BP, agreeing with the radiocarbon chronology for when penguins began to occupy Lagoon Island. A cooling period and glacial advance began at about 5000 BP and lasted for approximately 1000 years. Although this advance is not recorded in northern Marguerite Bay specifically, where the abandoned colonies are located, no calibrated penguin dates occur within that interval. Dates younger than 4000 BP occur within warming periods (the penguin optimum, 4000–3000 BP) or glacial

**Table II.** Alternate  $\Delta R$  values for calibrating three radiocarbon dates (in years BP) on penguin (*Pygoscelis* sp.) bone collagen from abandoned colonies on Lagoon Island, near Rothera Point. The sample lab number,  $\Delta R$  value, mean corrected date, and calibrated  $2\sigma$  range (95% confidence interval) are provided for each sample.

Sample/method	Corrected date	Calibrated range ( $2\sigma$ )
Beta 141907		
$\Delta R = 700 \pm 50$	$1210 \pm 60$	895–650
$\Delta R = 750 \pm 50$	$1160 \pm 60$	820–620
$\Delta R = 800 \pm 50$	$1110 \pm 60$	760–560
Beta 141911		
$\Delta R = 700 \pm 50$	$3540 \pm 70$	3585–3265
$\Delta R = 750 \pm 50$	$3490 \pm 70$	3530–3210
$\Delta R = 800 \pm 50$	$3440 \pm 70$	3460–3150
Beta 141916		
$\Delta R = 700 \pm 50$	$5380 \pm 70$	5900–5590
$\Delta R = 750 \pm 50$	$5330 \pm 70$	5870–5570
$\Delta R = 800 \pm 50$	$5280 \pm 70$	5770–5510

oscillations (3000 BP to present) and lack resolution for precise correlation with climatic events.

These data suggest that penguins were unable to occupy islands in the Rothera region until deglaciation exposed ice-free terrain at about 6000 BP. This first occupation lasted for approximately 1000 years, then ceased perhaps due to a glacial advance or increased snow deposition that again covered the islands. A warming trend at 4000 to 3000 BP can be correlated with re-occupation of Lagoon Island, after which glacial oscillations occur with no consistent correlation with the penguin occupation. The correspondence of these two independent records suggests that the  $\Delta R$  correction of  $700 \pm 50$  years and the calibration method is accurate. Moreover, the close association of three dates on squid (*Psychroteuthis glacialis*) with those from penguin bones from the same stratigraphical levels at Biscoe Point and Torgersen Island (Table I) indicates that this calibration method is appropriate for other krill-eating species in the Antarctic marine system. Thus, this method to correct for the marine-carbon reservoir effect is recommended for all radiocarbon dates of pygoscelid penguin remains in Antarctica.

The absence of abandoned colonies older than approximately 540 years (based on calibrated ranges, Table I) in the northern end of the Antarctic Peninsula remains an enigma. Tatur *et al.* (1997, p. 414) also noted a striking absence of sites older than 1300 years (based on surface guano) and hypothesized that these sites may have been eroded and destroyed by "solifluction and polygonal ground-forming processes." Only limited evidence indicates that penguins occupied King George Island in the mid Holocene (Bochenski 1985, Barsch & Mäusbacher 1986, Tatur *et al.* 1997, Sun *et al.* 2000). Moreover, a sediment core from Lake Bocckella, Hope Bay, has produced guano remains that suggest a large population of penguins colonized the surrounding terrain around 5550 BP (Zale 1994a, 1994b). It is possible that ornithogenic soils and abandoned sites predating 540 BP simply have not yet been located in the northern Antarctic Peninsula region, or that glaciation during the Little Ice Age (AD 1500–1850; Grove 1988) was sufficient to cover most islands with snow and ice, perhaps scouring and erasing any previous record of penguin occupation. The ice-core and oxygen isotope record from Siple station indicates that the Antarctic Peninsula did experience colder climates than today during the Little Ice Age, more so than east Antarctica (Mosley-Thompson *et al.* 1990).

The preservation of ornithogenic soils rich in bone and eggshell fragments of mid Holocene age at Rothera Point tends to falsify the suggestion that glacial scouring should account for the absence of old sites in the northern peninsula area. In addition, marine terraces and raised beaches on small islands near Rothera Point, including Lagoon and Anchorage Islands, have numerous polygonal features (Smith 1996, Emslie personal observation 2000) indicating that ground-forming processes have been active in this region. Freeze-thaw action in the pebbly ornithogenic soils over thousands of years has reduced most penguin bones preserved in these

sediments to small rounded fragments, but these, as well as eggshell fragments and other organic remains, are still abundant and well preserved. Thus, if abandoned colonies can survive glacial processes at Rothera, other explanations are needed for their absence in the northern peninsula. It is likely that a combination of factors that limit penguin distribution were responsible for the occupation and abandonment of specific areas. More detailed ground surveys and test excavations are warranted, especially on the high terraces and large ice-free regions on King George, Livingston, Seymour, James Ross, and Vega islands. These islands are considered to have the greatest potential for locating abandoned colonies and ornithogenic soils of early to middle Holocene age. Location of these sites would provide valuable information not only on the occupation history and diet of penguins, but also on the glacial and sea-level history of the northern Antarctic Peninsula and Weddell Sea region.

### Acknowledgements

This research was funded by Grant NAG5-8114 from the National Aeronautics and Space Administration. Research at Rothera station was made possible through a collaboration with the British Antarctic Survey. I thank A. Clarke, L. Peck, and P. Rose for their help and support during my visit; Jenny McDaniel assisted with fieldwork. C. Hjort kindly provided one unpublished radiocarbon date (UA 1033) from Devil Island and R.I. Lewis Smith provided three dates from Rothera Point and Lagoon Island. D. Hood and R. Hatfield, Beta Analytic, Inc, assisted in calibrating all radiocarbon dates reported here. The paper was improved with comments from C. Hjort, R.I. Lewis Smith and C. Baroni.

### References

- BARONI, C. & OROMBELLI, G. 1991. Holocene raised beaches at Terra Nova Bay, Victoria Land, Antarctica. *Quaternary Research*, **36**, 157–177.
- BARONI, C. & OROMBELLI, G. 1994. Abandoned penguin colonies as Holocene palaeoclimatic indicators in Antarctica. *Geology*, **22**, 23–26.
- BARSCH, D. & MÄUSBACHER, R. 1986. New data on relief development of the South Shetland Island, Antarctica. *International Science Review*, **11**, 211–218.
- BERKMAN, P.A. & FORMAN, S.L. 1996. Pre-bomb radiocarbon and the reservoir correction for calcareous marine species in the Southern Ocean. *Geophysical Research Letters*, **23**, 363–366.
- BERKMAN, P.A., ANDREWS, J.T., BJÖRCK, S., COLHOUN, E.A., EMSLIE, S.D., GOODWIN, I.D., HALL, B.L., HART, C.P., HIRAKAWA, K., IGARASHI, A., INGÓLFSSON, Ó., LÓPEZ-MARTINEZ, J., LYONS, W.B., MABIN, M.C.G., QUILTY, P.G., TAVIANI, M. & YOSHIDA, Y. 1998. Circum-Antarctic coastal environmental shifts during the late Quaternary reflected by emerged marine deposits. *Antarctic Science*, **10**, 345–362.
- BJÖRCK, S., HJORT, C., INGÓLFSSON, Ó. & SKOG, G. 1991. Radiocarbon dates from the Antarctic Peninsula region - problems and potential. *Quaternary Proceedings*, **1**, 55–65.
- BOCHENSKI, Z. 1985. Remains of subfossil birds from King George Island (South Shetland Islands). *Acta Zoologica Cracov*, **29**, 109–116.

- BRITISH ANTARCTIC SURVEY. 1989. *Proposed construction of airstrip at Rothera Point, Antarctica: final comprehensive environmental evaluation*. Cambridge: British Antarctic Survey, 57 pp.
- CAMPBELL, I.B. & CLARIDGE, G.G.C. 1987. *Antarctica: soils, weathering processes and environment*. Amsterdam: Elsevier, 355 pp.
- EMSLIE, S.D. 1995. Age and taphonomy of abandoned penguin colonies in the Antarctic Peninsula region. *Polar Record*, **31**, 409–418.
- EMSLIE, S.D., FRASER, W., SMITH, R.C. & WALKER, W. 1998. Abandoned penguin colonies and environmental change in the Palmer station area, Anvers Island, Antarctic Peninsula. *Antarctic Science*, **10**, 257–268.
- GORDON, J.E. & HARKNESS, D.D. 1992. Magnitude and geographic variation of the radiocarbon content in Antarctic marine life: implications for reservoir corrections in radiocarbon dating. *Quaternary Science Reviews*, **11**, 697–708.
- GROVE, J.M. 1988. *The Little Ice Age*. London: Methuen, 463 pp.
- HARKNESS, D.D. 1979. Radiocarbon dates from Antarctica. *British Antarctic Survey Bulletin*, No. 47, 43–59.
- HARRINGTON, H.J. & MCKELLAR, I.C. 1958. A radiocarbon date for penguin colonization of Cape Hallett, Antarctica. *New Zealand Journal of Geology and Geophysics*, **1**, 571–576.
- HEINE, J.C. & SPEIR, T.W. 1989. Ornithogenic soils of the Cape Bird Adélie penguin colonies, Antarctica. *Polar Biology*, **10**, 89–99.
- INGÓLFSSON, Ó., HJORT, C., BERKMAN, P.A., BJÖRCK, S., COLHOUN, E., GOODWIN, I.D., HALL, B., HIRAKAWA, K., MELLES, M., MÖLLER, P. & PRENTICE, M.L. 1998. Antarctic glacial history since the Last Glacial Maximum: an overview of the record on land. *Antarctic Science*, **10**, 326–344.
- LU, C.C. & WILLIAMS, R. 1994. Contribution to the biology of squid in the Prydz Bay region, Antarctica. *Antarctic Science*, **6**, 223–229.
- MOORS, P.J., SPEIR, T.W. & LYON, G.L. 1988. Soil analyses and <sup>13</sup>C/<sup>12</sup>C ratios identify sites of deserted rockhopper penguin colonies. *Auk*, **105**, 796–799.
- MOSLEY-THOMPSON, E., THOMPSON, L.G., GROOTES, P.M. & GUNDERSTRUP, N. 1990. Little Ice Age (Neoglacial) paleoenvironmental conditions at Siple station, Antarctica. *Annals of Glaciology*, **14**, 199–204.
- MYRCHA, A., PIETR, S.J. & TATUR, A. 1985. The role of pygoscelid penguin colonies in nutrient cycles at Admiralty Bay, King George Island. In SIEGFRIED, W.R., CONDY, P.R. & LAWS, R.M., eds. *Antarctic nutrient cycles and food webs*. New York: Springer, 156–162.
- SMITH, R.C., AINLEY, D., BAKER, K., DOMACK, E., EMSLIE, S., FRASER, B., KENNETT, J., LEVENTER, A., MOSLEY-THOMPSON, E., STAMMERJOHN, S. & VERNET, M. 1999. Marine ecosystem sensitivity to climate change. *Bioscience*, **49**, 393–404.
- SMITH, R.I.L. 1996. Terrestrial and freshwater biotic components of the western Antarctic Peninsula. *Antarctic Research Series*, **70**, 15–59.
- SPEIR, T.W. & COWLING, J.C. 1984. Ornithogenic soils of the Cape Bird Adélie penguin colonies, Antarctica. 1. Chemical properties. *Polar Biology*, **2**, 199–205.
- SPEIR, T.W. & HEINE, J.C. 1982. Penguin guano, decomposition and influence on soil development. *New Zealand Soil Bureau Record*, **83**, 1–39.
- SPEIR, T.W. & ROSS, D.J. 1984. Ornithogenic soils of the Cape Bird Adélie penguin colonies, Antarctica. 2. Ammonia evolution and enzyme activities. *Polar Biology*, **2**, 207–212.
- SPELLERBERG, I.F. 1970. Abandoned penguin colonies near Cape Royds, Ross Island, Antarctica and <sup>14</sup>C dating of penguin remains. *New Zealand Journal of Science*, **13**, 380–385.
- STONEHOUSE, B. 1970. Recent climatic change in Antarctica suggested from <sup>14</sup>C dating of penguin remains. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **7**, 341–343.
- STUIVER, M., PEARSON, G.W. & BRAZIUNAS, T. 1986. Radiocarbon age calibration of marine samples back to 9000 cal. yr BP. *Radiocarbon*, **28**, 980–1021.
- STUIVER, M. & REIMER, P.J. 1993. Extended <sup>14</sup>C data base and revised CALIB 3.0 <sup>14</sup>C age calibration program. *Radiocarbon*, **35**, 215–230.
- STUIVER, M., REIMER, P.J., BARD, E., BECK, J.W., BURR, G.S., HUGHEN, K.A., KROMER, B., MCCORMAC, G., VAN DER PLICHT, J. & SPURK, M. 1998. INTCAL98 radiocarbon age calibration, 24,000–0 cal BP. *Radiocarbon*, **40**, 1041–1083.
- SUN, L., XIE, Z. & ZHAO, J. 2000. A 3,000-year record of penguin populations. *Nature*, **407**, 858.
- SYROECHKOVSKY, E. 1959. The role of animals in the formation of primary soils under the conditions of circumpolar regions of the earth (Antarctica). *Zoologicheskyy Zhurnal*, **38**, 1770–1775.
- TATUR, A. 1987. Fluorine in ornithogenic soils and minerals on King George Island, West Antarctica. *Polish Polar Research*, **8**, 65–74.
- TATUR, A. 1989. Ornithogenic soils of the maritime Antarctic. *Polish Polar Research*, **10**, 481–532.
- TATUR, A. & MYRCHA, A. 1989. Soils and vegetation in abandoned penguin colonies (maritime Antarctic). *Proceedings NIPR Symposium on Polar Biology*, **2**, 181–189.
- TATUR, A., MYRCHA, A., FABISZEWSKI, J. & NIEGODZISZ, J. 1997. Formation of abandoned penguin colony ecosystems in maritime Antarctic. *Polar Biology*, **17**, 405–417.
- TEDROW, J.C.F. & UGOLINI, F.C. 1966. Antarctic soils. *Antarctic Research Series*, **8**, 161–177.
- UGOLINI, F.C. 1972. Ornithogenic soils of Antarctica. *Antarctic Research Series*, **20**, 181–193.
- ZALE, R. 1994a. Changes in size of the Hope Bay Adélie penguin colony as inferred from Lake Boeckella sediment. *Ecography*, **17**, 297–304.
- ZALE, R. 1994b. <sup>14</sup>C age corrections in Antarctic lake sediments inferred from geochemistry. *Radiocarbon*, **36**, 173–185.