

**Rapid Assessment and Monitoring of Coral Reef Habitats on the Florida Keys Reef Tract:
Quick Look Report: Summer 2002**

Principal Investigator:

Steven L. Miller, Center for Marine Science Research and NOAA's National Undersea Research Center, University of North Carolina at Wilmington, 515 Caribbean Drive, Key Largo, FL 33037, Tel: 305 451 0233, Fax: 305 453 9719, Email: millers@uncwil.edu

UNCW Project Team:

Mark Chiappone (Project Manager), Center for Marine Science Research, University of North Carolina at Wilmington, 515 Caribbean Drive, Key Largo, FL 33037, Email: chiapponem@uncwil.edu

Dione W. Swanson, Center for Marine Science Research, University of North Carolina at Wilmington, 515 Caribbean Drive, Key Largo, FL 33037 and Division of Marine Biology and Fisheries, Rosenstiel School of Marine and Atmospheric Science, University of Miami, 4600 Rickenbacker Causeway, Miami, FL 33149, Email: dswanson@rsmas.miami.edu

Additional 2002 Field Team:

Mark Vermeij, Cooperative Institute for Marine and Atmospheric Studies, NOAA/NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Drive, Miami, FL 33149 USA, Email: mark.vermeij@noaa.gov

Dave Eaken, Florida Marine Research Institute, South Florida Regional Laboratory, 2796 Overseas Highway, Suite 119, Marathon, FL 33050, Tel: 305 289 2330

Cooperating Institutions and Funding:

Florida Keys National Marine Sanctuary, Emerson Associates International, Rosenstiel School of Marine and Atmospheric Science-University of Miami, NOAA's National Marine Fisheries Service, National Undersea Research Center at the University of North Carolina at Wilmington, State of Florida.

Project Summary

This NURC/UNCW quick look report summarizes the accomplishments and summary findings from our summer 2002 benthic coral reef sampling in the Florida Keys. Our 2002 benthic sampling was partnered with a Keys-wide cruise of reef fishes and spiny lobster, focusing on hard-bottom and coral reef habitats from Biscayne National Park to the Tortugas region. The primary focus of the 2002 fieldwork was to obtain information on the community structure and condition of reef benthos at 15-21 m depth (fore reef) along the Florida Keys reef tract, to compare to similar studies carried out in 1995. The fieldwork also served to supplement existing data sets on the status of reef and hard-bottom communities in Biscayne National Park, the Florida Keys, and Tortugas region. The 2002 sampling represents the fifth consecutive year of large-scale surveys of hard-bottom and coral reef habitats conducted by NURC/UNCW.

In addition to the deeper fore reef sites, the 2002 work also included patch reef, hard-bottom, and reef terrace communities offshore of Biscayne Bay, the Florida Keys, and in the Dry Tortugas. The 2002 field effort complemented Keys-wide benthic surveys of 80 sites during 1999, 45 sites across the continental shelf in the lower Florida Keys region during 2000, 86 sites Keys-wide during 2001, and 60 sites sampled in the Dry Tortugas during 1999-2000. During 2002, we conducted 26 days of SCUBA surveys between May 30 and June 30. Sixty-four sites were sampled from Biscayne National Park to the Dry Tortugas. The deeper reef focus in 2002 was designed specifically to compare with results obtained in 1995 during a previous Keys-wide expedition. During 2002, three of the Sanctuary's 23 zones were visited, with ten additional sites surveyed just seaward of the zone boundaries in deeper water and habitats similar to sites surveyed during 1995. The entire field effort was supported by NURC/UNCW (project number 2002-14) extended operations aboard the M/V *Spree*. Benthic surveys by the project team were complemented by reef fish surveys conducted by personnel from RSMAS-University of Miami and NOAA-National Marine Fisheries Service.

Variables measured during 2002 included our core set of variables similar to what was sampled in previous years, namely coverage, species richness, gorgonian density, coral density and size, and coral condition. We were also able to include several variables (but not at all sites) that were sampled during the 2001 assessment: urchin density and size; density of anemones, corallimorpharians, opisthobranch mollusks, cleaner shrimps, spiny lobster, and arrow crabs; and *in situ* measurements of topographic complexity. All of the variables were collected underwater on pre-formatted plastic slates. We were also able to take several hundred underwater digital photographs to further develop our archive of Florida Keys reef and hard-bottom habitats and tools for taxonomic identification of benthic invertebrates.

Below we provide a brief narrative on the overall goals of the NURC/UNCW benthic assessment effort and a variable-by-variable summary of findings from the 2002 Keys-wide cruise. We also include a list of published and planned papers and other products developed from this effort to date. Several appended tables summarize site-level coverage, species richness, and density estimates for many of the variables measured, and four figures are included detailing the location of survey sites in Biscayne National Park, the Florida Keys, and the Tortugas region.

Sampling Goals and Objectives

The 2002 sampling of coral reef and hard-bottom habitats throughout the extent of the Florida Keys reef tract, including the Biscayne Bay and Dry Tortugas regions, complemented a multi-year effort dating back to 1999 (with pilot studies conducted in 1998) to assess shallow-water (< 21 m) coral reef and hard-bottom habitat types in the Sanctuary, including most of the Sanctuary no-take zones (Ecological Reserve, Sanctuary Preservation Areas, Research Only Areas) established in 1997. The goals of the NURC/UNCW zone monitoring effort are three-fold:

- To assess the community structure and condition of benthic communities at multiple spatial scales, with particular reference to the Sanctuary no-take zones. Variation is assessed among habitat types and among regions for particular habitat types.
- To assess potential changes in coral reef communities due to “no-take” protection from fishing within the Sanctuary zones, as well as from changes caused by larger-scale factors, such as geography and water quality.
- To provide fishery-independent reef fish surveys (conducted at the same time as the benthic surveys) with detailed habitat information, to facilitate modeling efforts to evaluate essential fishery habitat.

The 2002 Keys-wide cruise focused on the following tasks:

- To survey multiple coral reef and hard-bottom habitat types in the Biscayne Bay region of the Florida Reef tract, to complement surveys of similar habitat types in the upper, middle, and lower Keys during 1999-2001.
- To survey multiple coral reef and hard-bottom habitat types in the Dry Tortugas, to complement surveys of similar habitat types in the Dry Tortugas during 1999-2000.
- To survey deeper (15-21 m depth) low-relief spur and groove and low-relief hard-bottom habitats throughout the Florida Reef Tract, to provide a temporal comparison to similar surveys conducted during 1995 from Biscayne Bay to the Dry Tortugas.

Logistics and Methods (Tables 1-4: Figures 1-4)

A two-stage stratified random sampling design was used to select sites during 2002. A grid system constructed in a geographic information system (GIS) was used to overlay the existing habitat map of the Florida Keys. Sites or “blocks” 200 m x 200 m in dimension were used to randomly select sites from the following regional and habitat strata (Table 1):

- Biscayne Bay region: offshore patch reef, low-relief hard-bottom, and low-relief spur and groove
- Upper to Lower Keys: offshore patch reef, low-relief hard-bottom, and low-relief spur and groove
- Marquesas Keys: medium-profile reefs
- Dry Tortugas National Park: multiple reef and hard-bottom habitats
- Tortugas Bank: deep reef terraces

The 2002 effort included 14 sites east of Biscayne Bay (Figure 1), 24 sites from Key Largo to Key West (Figures 2 and 3), two sites south of the Marquesas Keys (Figure 3), and 24 sites in the Tortugas region (Figure 4). The 2002 sampling primarily focused on low-relief spur and groove and low-relief hard-bottom from 15-21 m depth (Table 2). These two habitat types were sampled during a similar Keys-wide expedition during 1995 and the 2002 surveys were thus designed to provide a temporal comparison to this earlier study. Three of the low-relief spur and groove sites sampled from Key Largo to Key West were located within no-take zones: Carysfort Reef SPA, Conch Reef Research Only, and Western Sambo Ecological Reserve (Table 2). Because many of the Sanctuary no-take zones do not extend beyond 12 m depth, low-relief spur and groove reefs were selected seaward of zone boundaries near ten existing zones: Elbow Reef, French Reef, Molasses Reef, Davis Reef, Alligator Reef, Tennessee Reef, Eastern Sambo, Eastern Dry Rocks, Rock Key, and Sand Key. One site or block was assigned to each zone and a total of 64 sites were surveyed between May 30 and June 30 (Table 2).

The 2002 sampling effort (64 sites) required 26 field days underwater from May 30 to June 30. Four days were lost to bad weather or other logistical issues. All 26 days were supported by NURC/UNCW extended operations aboard the M/V *Spree*. The field effort required approximately three or four dives a day by three to four divers. The 2002 sampling involved NURC/UNCW staff surveying the benthos, complemented by concurrent surveys of reef fishes by scientists from RSMAS-UM and NOAA/NMFS and lobster surveys by scientists from FDEP/FMRI. Table 3 summarizes the diving statistics for this year. Over 360 hours of surveys by NURC/UNCW and reef fish surveyors were required to complete the sampling.

The 2002 surveys addressed the same variables measured during 1999-2001, in addition to several variables added to the existing design during 2001 (Table 4). Briefly, pre-determined GPS points were used to locate the survey site or block. Four independent 15 m transects were deployed in each block, labeled in a numbered series from 1 to 4. The length of the transect was reduced from 25 m used in previous years because we optimized our sampling effort based on a statistical review of the existing data. Benthic coverage was estimated on all four transects and was determined every 15 cm to yield 100 points per transect. The number of species of stony corals, gorgonians, and sponges were determined on all four transects within a 0.5 m swath on either side of a 15 m transect (total survey area = 15 m^2 per transect). Gorgonian density was determined along two of the four transects within a 0.5 m swath on either side of the transect to the 8 m mark (total survey area = 8 m^2 per transect). Coral density, size, and condition were also determined along two of the four transects with the length of each swath fixed at 10 m (total survey area = 10 m^2 per transect). The coral condition measurements included identification of bleaching, disease, and an assessment of the extent to which interactions of coral and other taxa caused tissue damage or mortality. Juvenile corals (< 4 cm max. diameter) were assessed along two of the four transects by randomly sampling ten 0.68 m x 0.45 m quadrats along each transect (total survey area = 3.12 m^2 per transect). Urchin density and test diameter, as well as the density of incidental marine invertebrates were assessed on all four transects at selected sites. Topographic complexity was measured along the four transects to describe bottom slope, maximum vertical relief, and the coverage of different relief categories along 1.0 m x 15 m swaths.

Summary of Results

The summary results presented below describe the principal variables measured during 2002 and focus on regional differences in the low-relief spur and groove habitat type from the Biscayne Bay region to the Tortugas. This latter discussion will be the focus of a paper comparing the recent Keys-wide survey with a similar study conducted in 1995.

Benthic cover (Table 5)

Mean percent coverage data for scleractinian corals, fire coral (*Millepora* spp.), sponges, macroalgae, and algal turf are presented in Table 5. Patterns in coverage exhibited differences among the habitat types and broad regions surveyed. Overall, survey locations within the Biscayne region, across most of the habitat types, exhibited the lowest coral cover of the sites surveyed during 2002. Similar to earlier expeditions, offshore patch reefs exhibited some of the highest coral cover in the Biscayne and Florida Keys regions, ranging from 4% to nearly 28%. High-relief spur and groove (Bird Key Reef) and reef terrace habitats exhibited the highest coral cover in the Tortugas region and reef terraces exhibited the highest coral cover among all sites surveyed in 2002, ranging from 12% to 51%.

Offshore sampling locations in Biscayne and the Keys included three habitat types: low-relief spur and groove, low-relief hard-bottom, and patchy hard-bottom in sand. Not unexpectedly, coral cover offshore was greatest on low-relief spur and groove reefs, ranging from about 1% to nearly 15%. Surprisingly, coral cover did not tend to be greatest within the no-take zones. This contrasts with 2001 surveys of high-relief spur and groove reefs, in which sites within the no-take zones tended to have greater coral cover than adjacent reference sites. These results point to the value of sampling multiple habitat types, which often vary significantly even within the boundaries of individual Sanctuary no-take zones.

Sampling locations in the Dry Tortugas region included patch reefs or dead *Acropora cervicornis* mounds, low-relief hard-bottom, low-relief spur and groove, medium-profile reefs, high-relief spur and groove, reef knolls, and reef terraces. Reef terraces on the Tortugas Bank and on the western rim of Dry Tortugas National Park had the greatest coral cover measured among all habitat types during 2002. Algal cover in this habitat type was composed largely of brown foliose algae, especially *Lobophora variegata*.

Among the four regions surveyed in the low-relief spur and groove habitat, coral cover was greatest in the Dry Tortugas (1 site, 20.3%), followed by the middle Keys (mean of 7.4% among 5 sites), the lower Keys (6.8%), and the upper Keys (3%), the latter including sites east of Biscayne Bay. While coral cover varied the most among sites in the middle Keys (3.7% to 14.6%), we did not expect greater coral cover in this region relative to previous surveys in 1995. In fact, the pattern on deeper spur and groove reefs is opposite to that observed for shallower spur and groove reefs surveyed during 2001, in which reefs in the upper and lower Keys yielded the greatest coral cover. These results provide further support for broad geographic sampling across multiple habitat types and depths.

Species richness (Tables 6, 7)

Surveys of the number of species of stony corals, gorgonians, and sponges continued during the 2002 surveys. Table 6 lists the coral, gorgonian, and sponge species surveyed from all sites during 2002. A total of 46 coral species, 33 gorgonians, and 80 sponge species were found. The total numbers of species surveyed by site for these three invertebrate groups are summarized by habitat and region in Table 7. Similar to results from 2000 and 2001, patch reefs in the Keys and Biscayne regions typically yielded the greatest number of species of reef-building corals (Table 7). The number of sponges found on patch reefs in 2002 was similar or less than the number found on offshore fore reef areas.

Overall, coral species numbers surveyed in the low-relief spur and groove habitat were highest in the middle and lower Keys. No-take zones did not have significantly greater species numbers. Gorgonian species numbers were highest on patch reefs. On low-relief spur and groove reefs, gorgonian species richness was slightly higher in the lower Keys. Sponge species numbers were similar throughout the region, among habitats, and level of protection. In the Dry Tortugas, coral species numbers were similar among all habitat types with low-relief hard-bottom at the low end of the range. Both gorgonian and sponge species numbers were lowest in reef terrace habitats, but similar among all other habitats.

In the low-relief spur and groove habitat from east of Biscayne Bay to the lower Florida Keys, the mean number of coral species per site was greatest in the middle Keys (22.2), followed by the lower Keys (21.9) and the upper Keys (16.9). This is similar to findings from shallower spur and groove reefs surveyed during 2001, in which the upper Keys yielded the lowest species richness of corals. Similar to findings from 1999 and 2001, sponge species richness was, on average, greatest in the middle Keys (39.6 species per site), compared to the upper (37.8) and lower Keys (35.9). Gorgonian species richness did not vary significantly among the three regions within this particular habitat type, ranging from an average of 14.9 to 16.9 species per site.

Coral density, size and condition (Table 8)

Coral density, size, and condition measurements were made using methods similar to previous years. Table 8 lists the density of scleractinian corals by site. The total area surveyed during 2002 was approximately 1,280 m². Over 7,250 corals were counted and measured from the 64 sites, of which 2,098 or 29% were fire coral (*Millepora alcicornis*) and 5,161 or 71% were scleractinian corals.

Although scleractinian coral densities are highly variable, there were some patterns present. In the broad Biscayne and Keys area, scleractinian corals exhibited differences in density among habitat types and regional sectors. Coral density was highest on offshore patch reefs in the lower Keys. Densities were relatively similar among all other habitats in the lower, middle, and upper Keys, while density values in the Biscayne region were lower than the Keys for all habitat types sampled. In the Dry Tortugas, the highest coral densities were found on reef terraces, reef knolls, high-relief spur and groove (Bird Key reef), and low-relief spur and groove. Within the low-relief spur and groove habitat, mean coral densities within particular regions were greatest in the Dry Tortugas (1 site, 9.55 colonies/m²), followed by the middle Keys (5.49 colonies/m²). Size and condition data were not analyzed for this report.

Juvenile coral density (Table 8)

Surveys of juvenile coral species composition, density, and maximum diameter continued during 2002. Table 8 lists the density of juvenile scleractinian coral for each survey location. The highest juvenile coral densities were found on offshore patch reefs in the lower Keys. Juvenile coral densities were relatively similar, but variable within and among habitat types in the lower, middle, and upper Keys (excluding offshore patch reefs) and relatively higher than densities within and among habitats in the Biscayne area. The overall range of density values in the Dry Tortugas was similar to Keys-wide values, within and among habitat types. Medium-profile reefs exhibited the highest juvenile scleractinian coral densities overall. Within the low-relief spur and groove habitat among the upper, middle, and lower Keys, mean juvenile densities were greatest in the lower (7.1 juveniles/m²) and middle Keys (7.0 individuals per m²) and significantly lower in the upper Keys (4.7 juveniles/m²) for the 10 sites surveyed in this region.

Gorgonian density (Table 8)

Total gorgonian density for each site surveyed is listed in Table 8. From Biscayne to the Lower Keys, gorgonian densities were generally similar among habitats, regions, and management protection. The highest densities were recorded from offshore patch reefs in the lower Keys. Only nine sites in the Dry Tortugas included surveys for gorgonian density due to logistical constraints (three divers instead of four). Among the Tortugas sites, the highest gorgonian densities were found in the low-relief hard-bottom and high-relief spur and groove (Bird Key reef) habitats.

Urchin density and size (Table 9)

Surveys of urchin density were conducted at 56 of the 64 sites during 2002. Three species were encountered in transect surveys (Table 9). Similar to results from 1999-2001, all of the sampling locations yielded very low densities of urchins, particularly the long-spined sea urchin *Diadema antillarum*. However, we found some locations with relatively high densities of other species, particularly *Echinometra viridis*. Because densities remain low, patterns are difficult to discern, although we detected an abundance of *E. viridis* on offshore patch reefs compared to *Eucidaris tribuloides*, from Biscayne Bay to Key West.

Incidental invertebrates (Tables 10, 11, 12)

We assessed density patterns at 18 of the 64 sites surveyed for a variety of sessile and mobile invertebrate species during 2002 (Table 10). These surveys were not included at all sites for logistical reasons (three divers instead of four). In addition to anemones and corallimorpharians (Table 11), three species of shrimp symbionts, the polychaete *Hermodice carunculata*, the basket star *Astrophyton muricatum*, and crustaceans were counted within strip transect surveys at a number of sites (Table 12).

Plans for Use of the Data

We have made significant progress in manuscript development since January 2002. Below is a listing of manuscripts in press or published, those submitted for review, and those we intend to submit for publication by January 2003. We plan on using the 2002 data collected from the deeper fore reef to compare to a similar Keys-wide study conducted in 1995 of low-relief spur and groove habitats at 15-21 m depth.

Manuscripts published

1. Ault JS, Smith SG, Meester GA, Luo J, Bohnsack JA, Miller SL (2002) *Baseline multispecies coral reef fish stock assessment for the Dry Tortugas*. NOAA Technical Memorandum NMFS-SEFSC-487, 117 p
2. Ault JS, Smith SG, Meester GA, Luo J, Franklin EC, Bohnsack JA, Harper DE, McClellan DB, Miller SL, Swanson DW, Chiappone M (2002) Tortugas surveyed: Synoptic habitat and reef fish surveys support establishment of marine reserves in the Dry Tortugas, Florida, USA. *Reef Encounter* 31: 22-23
3. Chiappone M, Miller SL, Swanson DW, Ault JS, Smith SG (2001) Comparatively high densities of the long-spined sea urchin in the Dry Tortugas, Florida. *Coral Reefs* 20: 137-138
4. Chiappone M, Miller SL, Swanson DW (2001) *Condylactis gigantea* – A giant comes under pressure from the aquarium trade in Florida. *Reef Encounter* 30: 29-31
5. Chiappone M, Swanson DW, Miller SL (2002) Density, spatial distribution and size structure of sea urchins in coral reef and hard-bottom habitats of the Florida Keys. *Marine Ecology Progress Series* 235: 117-126
6. Chiappone M, Swanson DW, Miller SL, Smith SG (2002) Large-scale surveys on the Florida Reef Tract indicate poor recovery of the long-spined sea urchin *Diadema antillarum*. *Coral Reefs* 21: 155-159
7. Chiappone M, White A, Swanson DW, Miller SL (2002) Occurrence and biological impacts of fishing gear and other marine debris in the Florida Keys. *Marine Pollution Bulletin* 44: 597-604
8. Miller SL, Chiappone M, Swanson DW, Ault JS, Smith SG, Meester GA, Luo J, Franklin EC, Bohnsack JA, Harper DE, McClellan DB (2001) An extensive deep reef terrace on the Tortugas Bank, Florida Keys National Marine Sanctuary. *Coral Reefs* 20: 299-300

Manuscripts in press

9. Chiappone M, Dienes H, Swanson DW, Miller SL (In press) Density and gorgonian host-occupation patterns by flamingo tongue snails (*Cyphoma gibbosum*) in the Florida Keys. *Caribbean Journal of Science*
10. Dienes H, Chiappone M, Miller SL, Swanson DW (In press) Density and fore reef habitat utilization patterns of the lettuce sea slug (*Tridachia crispata*) in the Florida Keys. *Bulletin of Marine Science*
11. Dienes H, Chiappone M, Swanson DW, Miller SL (In press) Impacts of lost fishing gear on coral reef sessile invertebrates of the Florida Keys National Marine Sanctuary. *Biological Conservation*
12. Franklin EC, Ault JS, Smith SG, Luo J, Meester GA, Diaz GA, Chiappone M, Swanson DW, Miller SL, Bohnsack JA (In press) Coral reef benthic habitat classification in the Tortugas region, Florida. *Marine Geodesy*
13. Miller SL, Swanson DW, Chiappone M (In press) Multiple spatial scale assessment of coral reef and hard-bottom community structure in the Florida Keys National Marine Sanctuary. *Proceedings of the 9th International Coral Reef Symposium, Bali*

Manuscripts in review

14. Chiappone M, Swanson DW, Miller SL (In review) Density and habitat utilization patterns of anemones and corallimorpharians (Anthozoa, Zoantharia) in the Florida Keys National Marine Sanctuary. *Coral Reefs*
15. Chiappone M, Swanson DW, Miller SL (In review) Large-scale density patterns of anemones and corallimorpharians on offshore coral reef habitats in the Florida Keys. *Bulletin of Marine Science*
16. Dienes H, Chiappone M, Swanson DW, Miller SL (In review) Spatial distribution and density of fishing gear in Florida Keys National Marine Sanctuary no-take zones. *Coral Reefs*
17. Precht WF, Miller SL (2002) Ecological shifts along the Florida reef tract: Is the past the key to the future? In *The destruction of coral reef ecosystems: Paleoecological perspectives on the human role in a global crisis*. Aronson RB (ed), Springer Verlag, NY

Manuscripts in progress

18. Chiappone M, Swanson DW, Miller SL (In progress) Impacts to coral reef sessile invertebrates from lost lobster trap gear in the Florida Keys National Marine Sanctuary. *Symposium on Effects of Fishing Activities on Benthic Habitats, American Fisheries Society*
19. Chiappone M, Swanson DW, Miller SL (In progress) Spatial distribution and impacts of lost hook-and-line fishing gear to sessile invertebrates in the Florida Keys National Marine Sanctuary. *Symposium on Effects of Fishing Activities on Benthic Habitats, American Fisheries Society*
20. Chiappone M, Swanson DW, Miller SL, Sullivan-Sealey KM (In progress) A hierarchical structural classification of Florida Keys coral reef and hard-bottom habitats. *Aquatic Conservation: Marine and Freshwater Ecosystems*

21. Chiappone M, Swanson DW, Miller SL, White A, Dienes H (In progress) A rapid method for assessing topographic complexity of coral reef and hard-bottom habitats. *Journal of Experimental Marine Biology and Ecology*
22. Miller SL, Chiappone M, Swanson DW (In progress) Long-term dynamics of Florida Keys acroporid reefs: History and implications of a phase shift. *Coral Reefs*
23. Miller SL, Chiappone M, Swanson DW, Ault JS, Smith SG, Franklin EC (In progress) Design-based surveys of coral reef and hard-bottom habitats in Dry Tortugas National Park and the Tortugas Bank, Florida. *Ecological Applications*
24. Miller SL, Gittings S, Chiappone M, Causey B, Swanson DW, White A (In progress) Changes (1994-2000) to benthic cover on a deep coral reef in the Florida Keys. *Coral Reefs*
25. Smith SG, Swanson DW, Miller SL, Ault JS, Chiappone M (In progress) Sampling survey approaches for coral reef assessment and monitoring in the Florida Keys. *Coral Reefs*
26. Swanson DW, Chiappone M, Miller SL (In progress) Habitat and regional variations in coral species richness and coverage in the Florida Keys. *Coral Reefs*
27. Swanson DW, Chiappone M, Miller SL (In progress) Disease prevalence on reef-building corals in the Florida Keys National Marine Sanctuary. *Marine Ecology Progress Series*

Figure 1. Survey locations in the Biscayne Bay area during 2002.




Figure 2. Survey locations in the upper and middle Florida Keys regions during 2002.




Figure 3. Survey locations in the lower Florida Keys and Marquesas Keys areas during 2002.




Figure 4. Survey locations in the Tortugas region during 2002.




Table 1. Sampling effort by habitat type and region in the Florida Keys during May-June 2002. The survey effort included 27 sites in Biscayne National Park (BNP) and Dry Tortugas National Park (DTNP), 3 sites within Sanctuary no-take zones, and 10 sites just seaward of no-take zones.

Habitat type	Regional sector	Management type	No. of sites	Effort (%)
Offshore patch reef	Biscayne	Reference areas	2	3.1
	Lower Keys	Reference areas	2	3.1
Patch reef (Staghorn mound)	Dry Tortugas	DTNP	2	3.1
Reef knoll	Dry Tortugas	DTNP	3	4.7
High-relief spur and groove	Dry Tortugas	DTNP	1	1.6
Medium profile	Dry Tortugas	Reference areas	3	4.7
	Dry Tortugas	DTNP	1	1.6
Patchy hard-bottom in sand	Biscayne	BNP	2	3.1
Low-relief hard-bottom	Biscayne	BNP	6	9.4
	Upper Keys	Reference areas	2	3.1
	Upper Keys	No-take zones	1	1.6
	Dry Tortugas	Reference areas	4	6.3
	Dry Tortugas	DTNP	5	7.8
Low-relief spur and groove	Biscayne	BNP	4	6.3
	Upper Keys	Reference areas	5	7.8
	Upper Keys	No-take zones	1	1.6
	Middle Keys	Reference areas	4	6.3
	Middle Keys	No-take zones	1	1.6
	Lower Keys	Reference areas	7	10.9
	Lower Keys	No-take zones	1	1.6
Low-relief spur and groove	Dry Tortugas	DTNP	1	1.6
Reef terrace	Dry Tortugas	DTNP	1	1.6
	Dry Tortugas	Tortugas North ER	4	6.3
Seagrass matrix	Dry Tortugas	DTNP	1	1.6
Total			64	100.0

Table 2. Survey locations along the Florida Keys reef tract during May-June 2002. Sites are arranged from northeast to southwest within each habitat type. Sites within Sanctuary no-take zones are noted with a single asterisk (*) and those adjacent and seaward of zone boundaries are double asterisked (**).

Habitat type/site location	Region	Latitude	Longitude	Depth (m)
<i>Offshore patch reef</i>				
Inshore and SW of Pacific Reef	Biscayne	25.19.817	80.10.385	8.23-10.36
Inshore of Pacific Reef	Biscayne	25.21.095	80.09.551	10.06-10.97
West of W. Washerwoman Shoal	Lower Keys	24.32.579	80.36.076	5.18-7.01
West Washerwoman Shoal	Lower Keys	24.32.637	81.34.341	6.10-7.62
<i>Patch reef (Staghorn mound)</i>				
Dry Tortugas National Park	DTNP	24.36.537	82.55.831	9.45-10.36
Dry Tortugas National Park	DTNP	24.37.798	82.56.704	7.62-8.23
<i>Reef knoll</i>				
Dry Tortugas National Park	DTNP	24.37.378	82.49.468	14.02-17.98
Dry Tortugas National Park	DTNP	24.38.408	82.57.709	22.25-23.77
Dry Tortugas National Park	DTNP	24.41.616	82.54.714	10.97-11.89
<i>High-relief spur and groove</i>				
Bird Key Reef	Park	24.37.319	82.51.627	11.89-14.63
<i>Medium-profile reef</i>				
Coal Bin (East of Cosgrove Shoal)	Marquesas	24.27.014	82.08.089	16.76-17.37
Cosgrove Shoal	Marquesas	24.27.338	82.14.830	19.81-21.94
Pulaski Shoal (just outside DTNP)	Reference	24.43.367	82.47.006	17.07-17.37
Dry Tortugas National Park	DTNP	24.43.365	82.51.045	19.51-20.12
<i>Patchy hard-bottom in sand</i>				
Between Star & Triumph Reef	Biscayne	25.30.668	80.07.149	7.32-7.62
North of Fowey Rocks	Biscayne	25.37.442	80.05.600	10.67-11.58
<i>Low-relief hard-bottom</i>				
SW of Pacific Reef	Biscayne	25.18.272	80.09.772	11.89-11.89
Ajax Reef	Biscayne	25.24.093	80.07.796	6.10-7.32
North of Ajax Reef	Biscayne	25.24.914	80.07.580	7.01-8.84
Star Reef	Biscayne	25.31.448	80.06.142	11.28-12.19
Ledberry Reef	Biscayne	25.32.671	80.05.744	14.63-15.24
East of Biscayne Bay	Biscayne	25.41.200	80.05.746	8.84-10.36
Carysfort Reef*	Upper Keys	25.14.015	80.12.594	7.62-7.92
Seaward of Watson's Reef	Upper Keys	25.09.985	80.15.375	10.36-10.97
Between Molasses & French Reef	Upper Keys	25.01.438	80.21.732	10.97-11.89
S. DTNP (just outside park)	Reference	24.32.294	82.56.846	17.37-17.37
Dry Tortugas National Park	DTNP	24.33.690	82.56.353	14.33-14.94
Dry Tortugas National Park	DTNP	24.33.982	82.57.235	15.85-17.37
Dry Tortugas National Park	DTNP	24.35.383	82.56.541	12.80-13.72
Dry Tortugas National Park	DTNP	24.36.272	82.57.437	13.11-14.02
Dry Tortugas National Park	DTNP	24.40.434	82.55.023	9.14-10.36
DTNP (NW of Loggerhead Key)	Reference	24.40.772	82.55.278	15.24-16.15
Dry Tortugas National Park	DTNP	24.41.076	82.53.059	10.06-11.28
Tortugas Bank	Tortugas Bank	24.37.725	83.06.100	22.56-23.16

Table 2. continued.

Habitat type/site location	Region	Latitude	Longitude	Depth (m)
<i>Low-relief spur and groove</i>				
Offshore Pacific Reef	Biscayne	25.21.226	80.08.489	15.85-17.68
Between Star and Triumph Reef	Biscayne	25.30.334	80.06.348	10.97-12.50
SW of Brewster Reef	Biscayne	25.33.441	80.06.304	5.18-6.10
South of Fowey Rocks	Biscayne	25.34.699	80.05.519	15.54-19.20
Carysfort Reef SPA*	Upper Keys	25.12.371	80.12.780	16.76-18.90
Elbow Reef SPA**	Upper Keys	25.08.615	80.15.251	16.15-17.07
North Dixie Shoal	Upper Keys	25.05.032	80.18.337	14.63-16.46
Dixie Shoal	Upper Keys	25.04.232	80.18.995	14.63-15.85
French Reef**	Upper Keys	25.02.103	80.20.908	16.46-19.51
Molasses Reef**	Upper Keys	25.00.292	80.22.728	15.85-18.59
Conch Reef Research Only*	Middle Keys	24.56.903	80.27.250	14.33-15.54
Davis Reef**	Middle Keys	24.55.223	80.30.243	15.54-17.07
Alligator Reef**	Middle Keys	24.50.504	80.37.319	16.76-17.98
Between Tennessee & Alligator	Middle Keys	24.49.953	80.38.336	15.24-15.54
Offshore of Tennessee RO**	Middle Keys	24.45.331	80.45.021	15.24-16.76
East of Pelican Shoal	Lower Keys	24.30.274	81.36.000	11.89-13.41
Eastern Sambo**	Lower Keys	24.29.467	81.39.396	14.02-15.24
Western Sambo*	Lower Keys	24.28.771	81.42.240	14.94-15.85
Between W. Sambo & E. Dry Rocks	Lower Keys	24.28.113	81.46.068	17.37-18.59
Eastern Dry Rocks**	Lower Keys	24.27.386	81.50.750	15.54-17.68
Rock Key**	Lower Keys	24.27.188	81.51.320	13.41-16.15
Sand Key**	Lower Keys	24.27.025	81.53.069	14.63-15.85
SW of Western Dry Rocks	Lower Keys	24.26.636	81.59.849	14.93-16.46
Dry Tortugas National Park	DTNP	24.43.326	28.50.408	14.94-15.54
<i>Reef terrace</i>				
Loggerhead Forest	Park	24.39.558	82.56.005	16.46-17.37
Sherwood Forest	Tortugas Bank	24.41.274	83.04.010	23.16-23.77
Sherwood Forest	Tortugas Bank	24.42.052	83.01.899	24.99-26.21
Sherwood Forest	Tortugas Bank	24.42.275	83.02.671	21.64-22.25
Sherwood Forest	Tortugas Bank	24.42.469	83.03.055	23.77-24.38
<i>Seagrass matrix community</i>				
DTNP	Park	24.39.129	82.49.369	6.10-6.10

Table 3. SCUBA diving effort in the Florida Keys during May-June 2002. Reef fish surveys reported are for dives conducted in conjunction with the benthic surveys.

Diver	Affiliation	No. of dives	Depth (ft.)	Bottom time (hrs.)
<i>Benthic surveys</i>				
Steven Miller	CMSR/UNCW	37	26-86	49.58
Mark Chiappone	CMSR/UNCW	21	19-66	23.48
Dione Swanson	CMSR/UNCW	60	17-86	75.53
Mark Vermeij	RSMAS/UM	55	10-85	67.5
Dave Eaken	FDEP/FMRI	17	26-76	22.58
Subtotal		190	10-86	238.67
<i>Reef fish surveys</i>				
Steve Smith	RSMAS/UM	42	17-81	41.33
Mike Judge	NOAA/NMFS	21	30-79	21.67
Nicholas Farmer	RSMAS/UM	13	33-62	10.63
Mike Larkin	RSMAS/UM	22	28-65	17.77
Aaron Bartholomew	NOAA/NMFS	5	28-59	3.68
Lance Jordan	Nova Southeastern	12	22-70	11.58
Rob Waara	NPS/Virgin Islands	5	53-83	4.83
Brian Ettinger	Nova Southeastern	12	26-63	10.33
Subtotal		132	17-83	121.82
Total all divers		322	10-86	360.49

Table 4. Variables measured during 2002. Transects 15 m in length were used in all sites. Note that the width of the survey area along transects varied among variables.

Variable	Method	Factors assessed
Percent cover	Point-intercept along 4 transects (100 points/transect)	Percent cover
Species richness	1.0 m x 15 m swaths along 4 transects	Species presence & total number
Coral density and size	1.0 m x 10 m swaths along 2 transects	Density, size, condition
Juvenile coral density	Twenty 0.68 m x 0.45 m quadrats	Species composition, density & size
Gorgonian density	1.0 m x 8 m swaths along 2 transects	Density
Urchin density and size	1.0 m x 15 m swaths along 4 transects	Density, test diameter
Marine ornamentals	1.0 m x 15 m swaths along 4 transects	Density
Spiny lobster density	1.0 m x 15 m swaths along 4 transects	Density
Topography	1.0 m x 15 m swaths along 4 transects	Maximum relief & substratum slope

Table 5. Mean (1 SE) percent coverage of scleractinian corals, fire coral (*Millepora* spp.), sponges, macroalgae, and algal turf. Noted are sites within Sanctuary no-take zones (*) and those adjacent and seaward of the no-take zones (**). Data are based upon 100 points surveyed along each of four transects.

Habitat type/site location	Scleractinia	<i>Millepora</i>	Sponges	Macroalgae	Algal turf
<i>Offshore patch reef</i>					
Inshore and SW of Pacific Reef	3.8 (4.8)	1.3 (1.7)	0.3 (0.3)	35.9 (30.7)	21.2 (22.3)
Inshore of Pacific Reef	5.3 (6.6)	0.0 (0.0)	4.8 (6.0)	21.8 (22.7)	23.3 (23.8)
West Washerwoman Shoal	27.3 (26.4)	0.8 (1.0)	0.0 (0.0)	1.8 (2.3)	39.0 (31.7)
West of West Washerwoman	24.8 (24.8)	0.3 (0.3)	0.0 (0.0)	0.5 (0.7)	49.3 (33.3)
<i>Patch reef (Staghorn mound)</i>					
Dry Tortugas National Park	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	72.8 (26.4)	9.0 (10.9)
Dry Tortugas National Park	0.3 (0.3)	0.5 (0.7)	0.0 (0.0)	47.3 (33.2)	4.3 (5.4)
<i>Reef knoll</i>					
Dry Tortugas National Park	14.0 (24.1)	0.5 (1.0)	3.0 (5.8)	7.0 (13.0)	27.0 (39.4)
Dry Tortugas National Park	31.7 (32.5)	0.0 (0.0)	0.0 (0.0)	22.7 (26.3)	22.3 (26.0)
Dry Tortugas National Park	2.5 (3.2)	0.5 (0.7)	0.0 (0.0)	43.7 (32.8)	7.0 (8.7)
<i>High-relief spur and groove</i>					
Bird Key Reef	25.8 (25.5)	0.8 (1.0)	0.0 (0.0)	11.3 (13.3)	24.5 (24.7)
<i>Medium-profile reef</i>					
Coal Bin (East of Cosgrove Shoal)	5.0 (6.3)	0.0 (0.0)	0.0 (0.0)	56.5 (32.8)	21.3 (22.3)
Cosgrove Shoal	9.3 (11.2)	0.0 (0.0)	0.0 (0.0)	56.8 (32.7)	25.3 (25.2)
Dry Tortugas National Park	16.8 (18.6)	0.0 (0.0)	0.0 (0.0)	15.8 (17.7)	37.3 (31.2)
Pulaski Shoal	7.5 (13.9)	0.0 (0.0)	6.0 (11.3)	26.5 (39.0)	12.5 (21.9)
<i>Patchy hard-bottom in sand</i>					
Between Star & Triumph Reef	3.0 (3.9)	0.0 (0.0)	2.3 (2.9)	7.5 (9.3)	25.5 (25.3)
North of Fowey Rocks	0.0 (0.0)	0.8 (1.0)	0.3 (0.3)	6.3 (7.8)	49.3 (33.3)
<i>Low-relief hard-bottom</i>					
SW of Pacific Reef	0.8 (1.0)	0.5 (0.7)	0.5 (0.7)	21.3 (22.4)	50.7 (33.3)
Ajax Reef	1.5 (2.0)	0.5 (0.7)	3.8 (4.8)	34.8 (30.2)	37.8 (31.3)
North of Ajax Reef	0.8 (1.0)	0.0 (0.0)	2.0 (2.6)	30.8 (28.4)	45.0 (33.0)
Star Reef	0.5 (0.7)	0.3 (0.3)	0.8 (1.0)	21.5 (22.5)	48.5 (33.3)
Ledberry Reef	0.5 (0.7)	1.0 (1.3)	0.0 (0.0)	13.0 (15.1)	47.3 (33.2)
East of Biscayne Bay	2.0 (2.6)	0.5 (0.7)	0.8 (1.0)	8.5 (10.4)	53.5 (33.2)
Carysfort Reef*	2.5 (3.3)	0.5 (0.7)	1.2 (1.6)	36.0 (30.7)	34.1 (30.0)
Seaward of Watson's Reef	2.8 (3.6)	0.3 (0.3)	1.5 (2.0)	41.4 (32.4)	31.5 (28.8)
Between Molasses & French Reef	3.0 (3.9)	0.3 (0.3)	0.0 (0.0)	33.8 (29.8)	42.8 (32.6)
Southern DTNP (outside park)	2.5 (3.3)	0.3 (0.3)	0.0 (0.0)	42.0 (32.5)	12.0 (14.1)
Dry Tortugas National Park	3.3 (4.2)	0.5 (0.7)	0.0 (0.0)	42.3 (32.5)	13.3 (15.3)
Dry Tortugas National Park	3.5 (4.5)	0.3 (0.3)	0.0 (0.0)	44.0 (32.9)	17.5 (19.3)
Dry Tortugas National Park	4.0 (5.1)	1.3 (1.6)	0.5 (0.7)	35.0 (30.3)	37.8 (31.3)
Dry Tortugas National Park	8.3 (10.1)	1.3 (1.6)	0.0 (0.0)	33.8 (29.8)	6.0 (7.5)
Tortugas Bank	2.0 (2.6)	0.0 (0.0)	0.0 (0.0)	56.4 (32.8)	3.5 (4.5)
Dry Tortugas National Park	2.0 (2.6)	1.3 (1.6)	0.0 (0.0)	36.8 (31.0)	21.8 (22.7)
NW of Loggerhead Key	2.0 (2.6)	0.3 (0.3)	0.0 (0.0)	34.0 (29.9)	27.5 (26.6)
Dry Tortugas National Park	0.5 (0.7)	0.0 (0.0)	0.0 (0.0)	42.5 (32.6)	14.3 (16.3)

Table 5. continued.

Habitat type/site location	Scleractinia	Millepora	Sponges	Macroalgae	Algal turf
<i>Low-relief spur and groove</i>					
Offshore Pacific Reef	1.0 (1.3)	1.8 (2.3)	0.3 (0.3)	7.5 (9.3)	29.5 (27.7)
Between Star & Triumph Reef	2.5 (3.3)	0.0 (0.0)	1.5 (2.0)	32.3 (29.1)	35.8 (30.6)
SW of Brewster Reef	4.3 (5.4)	1.0 (1.3)	1.3 (1.6)	11.3 (13.3)	35.3 (30.4)
South of Fowey Rocks	3.2 (4.2)	0.6 (0.9)	0.6 (0.9)	21.0 (22.1)	28.5 (27.2)
Carysfort Reef SPA*	0.3 (0.3)	0.0 (0.0)	0.0 (0.0)	24.3 (24.5)	60.5 (31.9)
Elbow Reef SPA**	1.7 (2.3)	0.3 (0.3)	0.0 (0.0)	39.3 (31.8)	28.1 (26.9)
North Dixie Shoal	2.3 (2.9)	0.8 (1.0)	1.3 (1.6)	54.0 (33.1)	4.8 (6.0)
Dixie Shoal	6.5 (8.1)	0.5 (0.7)	0.0 (0.0)	49.1 (33.3)	26.2 (25.8)
French Reef**	4.0 (5.8)	0.3 (0.5)	0.3 (0.5)	51.7 (37.5)	10.3 (13.9)
Molasses Reef**	3.8 (4.8)	0.3 (0.3)	0.0 (0.0)	46.5 (33.2)	25.0 (25.0)
Conch Reef RO*	4.5 (5.7)	0.8 (1.0)	0.0 (0.0)	42.8 (32.6)	13.5 (15.6)
Davis Reef**	4.0 (5.1)	0.3 (0.3)	0.0 (0.0)	46.5 (33.2)	28.8 (27.3)
Alligator Reef**	3.7 (4.8)	1.7 (2.3)	0.0 (0.0)	19.6 (21.0)	20.3 (21.6)
Between Tennessee & Alligator	14.7 (16.7)	0.7 (1.0)	0.0 (0.0)	35.6 (30.6)	34.3 (30.0)
Offshore of Tennessee RO**	10.3 (12.3)	0.0 (0.0)	0.0 (0.0)	41.5 (32.4)	38.0 (31.4)
East of Pelican Shoal	0.8 (1.0)	1.0 (1.3)	0.0 (0.0)	37.2 (31.1)	16.2 (18.1)
Eastern Sambo Reef**	7.8 (9.5)	0.0 (0.0)	0.0 (0.0)	39.0 (31.7)	21.0 (22.1)
Western Sambo Reef*	10.3 (12.3)	0.0 (0.0)	0.0 (0.0)	42.0 (32.5)	19.8 (21.1)
Between W. Sambo & E. Dry Rocks	13.3 (15.3)	0.3 (0.3)	0.0 (0.0)	50.0 (33.3)	11.0 (13.1)
Eastern Dry Rocks**	7.5 (9.3)	0.5 (0.7)	0.0 (0.0)	45.8 (33.1)	38.0 (31.4)
Rock Key**	6.5 (8.1)	0.3 (0.3)	0.0 (0.0)	34.8 (30.2)	51.0 (33.3)
Sand Key**	3.3 (4.2)	0.0 (0.0)	0.0 (0.0)	63.0 (31.1)	27.3 (26.4)
SW of Western Dry Rocks	4.8 (6.0)	0.5 (0.7)	0.0 (0.0)	45.3 (33.0)	37.5 (31.3)
Dry Tortugas National Park	20.3 (21.5)	1.3 (1.6)	0.0 (0.0)	38.0 (31.4)	24.8 (24.8)
<i>Reef terrace</i>					
Loggerhead Forest	50.5 (33.3)	0.8 (1.0)	0.0 (0.0)	27.5 (26.6)	10.3 (12.3)
Sherwood Forest	29.8 (27.9)	0.3 (0.3)	0.0 (0.0)	35.0 (30.1)	0.5 (0.7)
Sherwood Forest	32.7 (33.0)	0.0 (0.0)	0.0 (0.0)	32.9 (33.1)	0.5 (0.8)
Sherwood Forest	24.0 (24.3)	0.0 (0.0)	0.0 (0.0)	41.3 (32.3)	10.0 (12.0)
Sherwood Forest	11.5 (13.6)	0.3 (0.3)	0.0 (0.0)	56.9 (32.7)	12.5 (14.6)

Table 6. Coral, gorgonian, and sponge species surveyed during May-June 2002. Listed are only those species recorded within four 15 m² plots per site.

Stony corals	Gorgonians	Sponges	
<i>Acropora cervicornis</i>	<i>Briareum asbestinum</i>	<i>Adocia</i> sp.	<i>Pandaros acanthifolium</i>
<i>Agaricia agaricites</i>	<i>Erythrop. caribaeorum</i>	<i>A. carbonifera</i>	<i>Pseudoaxinella lunaecharta</i>
<i>A. fragilis</i>	<i>Eunicea calyculata</i>	<i>Agelas clathrodes</i>	<i>Pseudoceratina crassa</i>
<i>A. grahamae</i>	<i>E. fusca</i>	<i>A. conifera</i>	<i>Ptilocaulis</i> sp.
<i>A. humilis</i>	<i>E. laciniata</i>	<i>A. schmidti</i>	<i>Rhaphidophlus juniperinis</i>
<i>A. lamarcki</i>	<i>E. mammosa</i>	<i>A. wiedenmayeri</i>	<i>R. venosus</i>
<i>Colpophyllia natans</i>	<i>E. palmeri</i>	<i>Amphimedon compressa</i>	<i>Siphon. Coralliphagum</i>
<i>Dichocoenia stokesi</i>	<i>E. succinea</i>	<i>A. viridis</i>	<i>S. siphonum</i>
<i>Diploria clivosa</i>	<i>E. tourneforti</i>	<i>Anthosigmella varians</i>	<i>Spheciospongia vesparium</i>
<i>D. labyrinthiformis</i>	<i>Gorgonia ventalina</i>	<i>Aplysina archeri</i>	<i>Spinoseella tenerrima</i>
<i>D. strigosa</i>	<i>Iciliogorgia schrammi</i>	<i>A. cauliformis</i>	<i>Spirastrella coccinea</i>
<i>Eusmilia fastigiata</i>	<i>Muricea atlantica</i>	<i>A. fistularis</i>	<i>Spongia</i> sp.
<i>Favia fragum</i>	<i>M. elongata</i>	<i>A. fulva</i>	<i>Strongylacidon</i> sp.
<i>Isophyllastrea rigida</i>	<i>M. muricata</i>	<i>A. lacunosa</i>	<i>Tedania ignis</i>
<i>Isophyllia sinuosa</i>	<i>M. pinnata</i>	<i>Callyspongia plicifera</i>	<i>Tethya crypta</i>
<i>Leptoseris cucullata</i>	<i>Muriceopsis flavida</i>	<i>C. vaginalis</i>	<i>Ulosa ruetzleri</i>
<i>Madracis carmabi</i>	<i>Plexaura flexuosa</i>	<i>Chondrilla nucula</i>	Unknown blue tube #1
<i>M. decactis</i>	<i>P. homomalla</i>	<i>Cinachyra</i> sp.	Unknown blue tube #2
<i>M. formosa</i>	<i>Plexaurella dichotoma</i>	<i>Clathria</i> sp.	Unknown brown <i>Cliona</i>
<i>M. mirabilis</i>	<i>P. grisea</i>	<i>Clathrina canariensis</i>	Unknown brown encrusting
<i>M. senaria</i>	<i>P. nutans</i>	<i>Cliona</i> sp.	Unknown brown lumpy
<i>Manicina areolata</i>	<i>P. pumila</i>	<i>C. deletrix</i>	Unknown brown smooth
<i>Meandrina meandrites</i>	<i>Pseudoplexaura crucis</i>	<i>C. langae</i>	Unknown brown tube
<i>Millepora alcicornis</i>	<i>P. flagellosa</i>	<i>Cribochalina vasculum</i>	Unknown carmine red
<i>M. complanata</i>	<i>P. porosa</i>	<i>Diplastrella megastellata</i>	Unknown encrusting
<i>Montastraea annularis</i>	<i>P. wagnaari</i>	<i>Dysidea etheria</i>	Unknown mauve lumpy
<i>M. cavernosa</i>	<i>Pseudopt. acerosa</i>	<i>Ectyoplasia ferox</i>	Unknown orange encrusting
<i>M. faveolata</i>	<i>P. americana</i>	<i>Erylus formosus</i>	Unknown red encrusting
<i>M. franksi</i>	<i>P. bipinnata</i>	<i>Geodia neptuna</i>	Unknown red lumpy
<i>Mussa angulosa</i>	<i>P. rigida</i>	<i>Haliclona hogarthi</i>	Unknown red squishy
<i>Mycetophyllia aliciae</i>	<i>Pterogorgia anceps</i>	<i>Halisarca</i> sp.	<i>Verongula gigantea</i>
<i>M. danaana</i>	<i>P. citrina</i>	<i>Holapsamma helwigi</i>	<i>V. rigida</i>
<i>M. ferox</i>	<i>P. guadalupensis</i>	<i>Iotrochota birotulata</i>	<i>V. reisiwigi</i>
<i>M. lamarckiana</i>		<i>Ircinia campana</i>	<i>Xestospongia muta</i>
<i>Oculina diffusa</i>		<i>Ircinia felix</i>	
<i>Porites astreoides</i>		<i>Ircinia strobilina</i>	
<i>P. branneri</i>		<i>Monanchora barbadensis</i>	
<i>P. colonensis</i>		<i>Monanchora unguifera</i>	
<i>P. porites divaricata</i>		<i>Mycale laevis</i>	
<i>P. porites furcata</i>		<i>Mycale</i> sp.	
<i>P. porites porites</i>		<i>Myrmekioderma</i> sp.	
<i>Scolymia</i> sp.		<i>Neofibularia notilangere</i>	
<i>Siderastrea radians</i>		<i>Niphates amorpha</i>	
<i>S. siderea</i>		<i>N. digitalis</i>	
<i>Solenastrea bournoni</i>		<i>N. erecta</i>	
<i>Steph. michelinii</i>		<i>Oligoceras hemorrhages</i>	

Table 7. Total number of species per site for stony corals, gorgonians, and sponges. Data are based upon four 15 m² surveys conducted at each site (60 m²). Noted are sites within Sanctuary no-take zones (*) and those adjacent and seaward of the no-take zones (**). ns = no survey conducted.

Habitat type/site location	Region	Stony corals	Gorgonians	Sponges
<i>Offshore patch reef</i>				
Inshore and SW of Pacific Reef	Biscayne	21	21	41
Inshore of Pacific Reef	Biscayne	20	21	42
West Washerwoman Shoal	Lower Keys	28	24	28
West of West Washerwoman	Lower Keys	28	21	33
<i>Patch reef (Staghorn mound)</i>				
Dry Tortugas National Park	Tortugas	5	ns	ns
Dry Tortugas National Park	Tortugas	4	ns	ns
<i>Reef knoll</i>				
Dry Tortugas National Park	Tortugas	23	15	39
Dry Tortugas National Park	Tortugas	17	ns	ns
Dry Tortugas National Park	Tortugas	16	ns	ns
<i>High-relief spur and groove</i>				
Bird Key Reef	Tortugas	29	22	43
<i>Medium-profile reef</i>				
Coal Bin (East of Cosgrove Shoal)	Marquesas	28	17	40
Cosgrove Shoal	Marquesas	20	14	35
Dry Tortugas National Park	Tortugas	20	ns	ns
Pulaski Shoal (just outside DTNP)	Tortugas	25	11	43
<i>Patchy hard bottom in sand</i>				
Between Star & Triumph Reef	Biscayne	13	23	40
North of Fowey Rocks	Biscayne	14	14	30
<i>Low-relief hard-bottom</i>				
SW of Pacific Reef	Biscayne	17	20	37
Ajax Reef	Biscayne	18	18	17
North of Ajax Reef	Biscayne	14	16	25
Star Reef	Biscayne	17	18	36
Ledberry Reef	Biscayne	14	12	35
East of Biscayne Bay	Biscayne	18	14	37
Carysfort Reef SPA*	Upper Keys	22	16	27
Seaward of Watson's Reef	Upper Keys	16	18	44
Between Molasses & French Reef	Upper Keys	19	15	38
Southern DTNP (just outside park)	Tortugas	16	20	43
Dry Tortugas National Park	Tortugas	21	23	38
Dry Tortugas National Park	Tortugas	15	21	42
Dry Tortugas National Park	Tortugas	25	22	48
Dry Tortugas National Park	Tortugas	18	ns	7
Dry Tortugas National Park	Tortugas	16	25	32
NW of Loggerhead Key	Tortugas	16	27	42
Dry Tortugas National Park	Tortugas	13	ns	ns
Tortugas Bank	Tortugas	19	13	35

Table 7. continued.

Habitat type/site location	Region	Stony corals	Gorgonians	Sponges
<i>Low-relief spur and groove</i>				
Offshore Pacific Reef	Biscayne	18	18	37
Between Star & Triumph Reef	Biscayne	18	18	43
SW of Brewster Reef	Biscayne	16	18	33
South of Fowey Rocks	Biscayne	17	7	37
Carysfort Reef SPA*	Upper Keys	16	13	38
Elbow Reef SPA**	Upper Keys	19	17	42
North Dixie Shoal	Upper Keys	18	13	31
Dixie Shoal	Upper Keys	19	18	39
French Reef**	Upper Keys	14	10	32
Molasses Reef**	Upper Keys	14	17	46
Conch Reef Research Only*	Middle Keys	19	12	39
Davis Reef**	Middle Keys	21	15	42
Alligator Reef**	Middle Keys	21	20	46
Between Tennessee & Alligator	Middle Keys	27	21	35
Offshore of Tennessee RO**	Middle Keys	23	16	36
East of Pelican Shoal	Lower Keys	16	24	34
Eastern Sambo**	Lower Keys	18	16	37
Western Sambo*	Lower Keys	27	10	32
Between W. Sambo & E. Dry Rocks	Lower Keys	22	9	22
Eastern Dry Rocks**	Lower Keys	23	18	39
Rock Key**	Lower Keys	22	19	41
Sand Key**	Lower Keys	23	20	38
SW of Western Dry Rocks	Lower Keys	24	19	44
Dry Tortugas National Park	Tortugas	30	14	40
<i>Reef terrace</i>				
Loggerhead Forest	Tortugas	28	7	25
Sherwood Forest	Tortugas	25	10	24
Sherwood Forest	Tortugas	25	9	31
Sherwood Forest	Tortugas	26	9	25
Sherwood Forest	Tortugas	19	ns	ns

Table 8. Mean (1 SE) density (no. colonies/m²) of scleractinian corals, juvenile scleractinian corals (< 4 cm max. diameter), and gorgonians. Noted are sites within Sanctuary no-take zones (*) and sites adjacent and seaward of no-take zone boundaries (**). ns = no survey conducted.

Habitat type/site location	Scleractinian corals	SE	Juvenile corals	SE	Gorgonians	SE
<i>Offshore patch reef</i>						
Inshore and SW of Pacific Reef	3.15	0.01	3.21	0.00	19.88	0.03
Inshore of Pacific Reef	4.25	0.61	5.45	0.21	11.00	0.78
West Washerwoman Shoal	13.65	1.13	16.51	37.44	27.88	12.50
West of West Washerwoman	13.75	0.85	13.14	10.07	11.44	9.57
<i>Patch reef (Staghorn mound)</i>						
Dry Tortugas National Park	0.30	0.02	0.16	0.05	ns	ns
Dry Tortugas National Park	0.00	0.00	0.00	0.00	ns	ns
<i>Reef knoll</i>						
Dry Tortugas National Park	7.55	1.45	5.13	0.82	ns	ns
Dry Tortugas National Park	8.15	3.13	4.41	0.00	ns	ns
Dry Tortugas National Park	1.60	0.02	2.40	0.05	ns	ns
<i>High-relief spur and groove</i>						
Bird Key Reef	8.25	0.05	7.05	0.21	31.92	0.35
<i>Medium-profile reef</i>						
Coal Bin (East of Cosgrove Shoal)	4.65	0.41	8.81	6.22	23.00	2.72
Cosgrove Shoal	4.60	0.02	6.41	7.40	12.88	4.50
Dry Tortugas National Park	7.20	8.82	6.09	1.85	ns	ns
Pulaski Shoal (just outside park)	5.30	0.00	7.21	4.16	ns	ns
<i>Patchy hard bottom in sand</i>						
Between Star & Triumph Reef	2.35	0.13	4.81	0.82	10.69	21.95
North of Fowey Rocks	1.70	1.28	5.61	1.28	10.88	0.28
<i>Low-relief hard-bottom</i>						
SW of Pacific Reef	2.15	0.61	3.85	3.29	17.10	27.38
Ajax Reef	2.30	0.32	1.60	0.00	14.81	0.20
North of Ajax Reef	1.40	0.08	5.45	1.85	10.94	0.95
Star Reef	1.20	0.02	4.65	0.05	11.00	4.50
Ledberry Reef	0.95	0.13	3.53	5.14	15.06	0.07
East of Biscayne Bay	3.10	0.18	6.25	2.52	8.94	8.51
Carysfort Reef SPA*	7.15	1.13	9.13	4.16	5.38	4.50
Seaward of Watson's Reef	3.95	3.65	10.74	4.16	16.13	78.13
Between Molasses & French Reef	2.35	0.41	3.53	0.82	10.13	0.13
Southern DTNP (just outside park)	1.95	0.13	3.85	0.82	19.17	0.22
Dry Tortugas National Park	1.40	0.08	4.43	0.96	25.33	26.89
Dry Tortugas National Park	2.30	0.32	5.13	0.00	26.83	2.72
Dry Tortugas National Park	3.90	1.28	6.89	0.46	21.94	35.07
Dry Tortugas National Park	2.05	0.25	4.17	0.21	ns	ns
Tortugas Bank	2.05	0.25	5.93	4.16	ns	ns
Dry Tortugas National Park	0.90	0.00	2.40	2.52	27.85	0.13
NW of Loggerhead Key	0.90	0.18	0.96	0.00	ns	ns
Dry Tortugas National Park	1.60	0.72	2.56	0.00	ns	ns

Table 8. continued.

Habitat type/site location	Scleractinia	SE	Juvenile corals	SE	Gorgonians	SE
<i>Low-relief spur and groove</i>						
Offshore Pacific Reef	1.95	0.05	3.69	1.28	9.56	20.32
SW of Brewster Reef	5.00	2.00	3.04	4.16	27.19	3.45
South of Fowey Rocks	3.95	2.21	4.81	5.14	4.69	3.45
Between Star & Triumph Reef	1.80	0.18	4.49	3.29	14.06	2.26
Carysfort Reef SPA*	1.20	0.02	4.33	0.46	9.75	0.28
Elbow Reef SPA**	3.35	0.85	7.37	5.14	10.56	0.01
North Dixie Shoal	5.40	0.50	4.81	3.29	6.13	0.03
Dixie Shoal	5.25	0.25	5.93	8.68	10.56	2.26
French Reef **	2.95	3.65	4.49	0.21	19.81	4.88
Molasses Reef**	3.55	0.25	4.01	4.16	10.25	3.78
Conch Reef Research Only*	3.95	1.13	10.26	0.82	8.19	0.95
Davis Reef**	3.95	0.01	6.57	6.22	ns	ns
Alligator Reef**	2.10	2.00	3.85	3.29	ns	ns
Between Tennessee & Alligator	11.31	0.20	10.42	27.17	ns	ns
Offshore of Tennessee RO**	6.13	0.78	3.69	6.22	ns	ns
East of Pelican Shoal	2.85	0.61	10.90	1.85	15.13	0.00
Eastern Sambo**	4.75	0.85	6.57	1.28	ns	ns
Western Sambo*	4.50	0.50	3.77	0.12	ns	ns
Between W. Sambo & E. Dry Rocks	5.48	3.25	ns	ns	ns	ns
Eastern Dry Rocks**	5.50	0.50	3.69	6.22	ns	ns
Rock Key**	3.80	2.00	9.46	1.28	ns	ns
Sand Key**	2.75	1.81	6.53	2.34	ns	ns
SW of Western Dry Rocks	3.80	0.72	7.85	4.16	ns	ns
Dry Tortugas National Park	9.55	0.25	8.01	1.85	13.08	3.13
<i>Reef terrace</i>						
Loggerhead Forest	5.35	1.13	1.32	0.93	ns	ns
Sherwood Forest	7.90	0.72	3.04	6.22	ns	ns
Sherwood Forest	7.75	0.05	ns	ns	ns	ns
Sherwood Forest	7.20	0.08	6.13	12.74	ns	ns
Sherwood Forest	4.70	0.72	2.75	0.07	ns	ns

Table 9. Mean (1 SE) densities of urchins (no. individuals per m²) from surveys of four 15 m² plots per site. Noted are sites within Sanctuary no-take zones (*) and those adjacent and seaward of zone boundaries (**). ns = no survey conducted.

Habitat type/site location	<i>Diadema antillarum</i>		<i>Eucidaris tribuloides</i>		<i>Echinometra viridis</i>	
	Mean	SE	Mean	SE	Mean	SE
<i>Offshore patch reef</i>						
Inshore of Pacific Reef	0	0	0.017	0.001	0	0
West Washerwoman Shoal	0	0	0	0	0.711	0.099
West of West Washerwoman	0.017	0.001	0.017	0.001	1.033	0.336
<i>Patch reef (Staghorn mound)</i>						
Dry Tortugas National Park	0	0	0	0	0	0
<i>Reef knoll</i>						
Dry Tortugas National Park	0.017	0.001	0	0	0.150	0.007
<i>High-relief spur and groove</i>						
Bird Key Reef	0	0	0	0	0.217	0.046
<i>Medium-profile reef</i>						
Coal Bin (East of Cosgrove Shoal)	0	0	0	0	0	0
Cosgrove Shoal	0	0	0	0	0.017	0.001
Pulaski Shoal (just outside DTNP)	0	0	0	0	0	0
<i>Patchy hard-bottom in sand</i>						
Between Star & Triumph Reef	0.050	0.004	0	0	0.033	0.001
North of Fowey Rocks	0	0	0	0	0	0
<i>Low-relief hard-bottom</i>						
SW of Pacific Reef	0	0	0	0	0	0
Ajax Reef	0	0	0.017	0.001	0.017	0.001
North of Ajax Reef	0	0	0	0	0	0
Star Reef	0	0	0	0	0	0
Ledberry Reef	0	0	0	0	0	0
East of Biscayne Bay	0	0	0	0	0	0
Carysfort Reef SPA*	0	0	0	0	0	0
Seaward of Watson's Reef	0	0	0	0	0.050	0.010
Between Molasses & French Reef	0	0	0.017	0.001	0.017	0.001
Southern DTNP (just outside park)	0	0	0.017	0.001	0	0
Dry Tortugas National Park	0	0	0.017	0.001	0	0
Dry Tortugas National Park	0	0	0	0	0	0
Dry Tortugas National Park	0	0	0	0	0.017	0.001
Tortugas Bank	0	0	0	0	0	0
Dry Tortugas National Park	0.022	0.001	0	0	0	0
NW of Loggerhead Key	0	0	0	0	0	0

Table 9. continued.

Habitat type/site location	<i>Diadema antillarum</i>		<i>Eucidaris tribuloides</i>		<i>Echinometra viridis</i>	
	Mean	SE	Mean	SE	Mean	SE
<i>Low- relief spur and groove</i>						
Offshore Pacific Reef	0	0	0	0	0	0
Between Star & Triumph Reef	0.017	0.001	0	0	0	0
SW of Brewster Reef	0	0	0.017	0.001	0	0
South of Fowey Rocks	0	0	0	0	0	0
Carysfort Reef SPA*	0	0	0	0	0	0
Elbow Reef SPA**	0	0	0	0	0	0
North Dixie Shoal	0	0	0	0	0	0
Dixie Shoal	0	0	0	0	0	0
French Reef**	0	0	0	0	0	0
Molasses Reef**	0	0	0	0	0	0
Conch Reef Research Only*	0	0	0	0	0	0
Davis Reef**	0	0	0	0	0	0
Alligator Reef**	0	0	0	0	0	0
Between Tennessee & Alligator	0	0	0.017	0.001	0	0
Offshore of Tennessee RO**	0.022	0.001	0	0	0	0
East of Pelican Shoal	0	0	0	0	0	0
Eastern Sambo Reef**	0	0	0	0	0	0
Western Sambo Reef*	0	0	0	0	0	0
Between W. Sambo & E. Dry Rocks	0	0	0	0	0	0
Eastern Dry Rocks	0	0	0	0	0	0
Rock Key**	0	0	0	0	0.017	0.001
Sand Key**	0	0	0	0	0	0
SW of Western Dry Rocks	0	0	0.022	0.001	0	0
Dry Tortugas National Park	0	0	0	0	0	0
<i>Reef terrace</i>						
Loggerhead Forest	0	0	0	0	0	0
Sherwood Forest	0	0	0	0	0	0
Sherwood Forest	0.022	0.001	0	0	0	0
Sherwood Forest	0	0	0	0	0.017	0.001
Sherwood Forest	0	0	0	0	0.017	0.001

Table 10. Incidental marine invertebrates surveyed for density during 2002. Listed are only those species recorded within four 15 m² plots per site.

Invertebrate group	Species name	Common name
Anemones	<i>Bartholomea annulata</i>	Ringed anemone
	<i>Condylactis gigantea</i>	Pink-tipped anemone
	<i>Lebrunia danae</i>	
Corallimorpharians	<i>Ricordea florida</i>	Florida corallimorph
Polychaetes	<i>Hermodice carunculata</i>	Fireworm
Echinoderms	<i>Astrophyton muricatum</i>	Basket star
	<i>Diadema antillarum</i>	Long-spined sea urchin
	<i>Echinometra viridis</i>	Reef urchin
	<i>Eucidaris tribuloides</i>	Red pencil urchin
Crustaceans	<i>Panulirus argus</i>	Spiny lobster
	<i>Periclemenes pedersoni</i>	Pederson's cleaner shrimp
	<i>Stenorhyncus seticornis</i>	Arrow crab
	<i>Stenopus hispidus</i>	Red-banded coral shrimp

Table 11. Mean (1 SE) densities (no. individuals per m²) of anemones and corallimorpharians from surveys of four 15 m² plots per survey location. Sites within Sanctuary no-take zones are noted with a single asterisk (*) and sites adjacent and seaward of zone boundaries are double asterisked (**). Listed are only the sites surveyed and species recorded within four 15 m² plots per site.

Habitat type/site location	<i>Bartholomea annulata</i>	<i>Condylactis gigantea</i>	<i>Lebrunia danae</i>	<i>Ricordea florida</i>
<i>Offshore patch reef</i>				
Inshore of Pacific Reef	0.033 (0.004)	0.017 (0.001)	0.017 (0.001)	0
<i>Patchy hard-bottom in sand</i>				
Between Star & Triumph Reef	0.033 (0.001)	0.033 (0.001)	0	0.150 (0.022)
North of Fowey Rocks	0.017 (0.001)	0	0	0
<i>Low-relief hard bottom</i>				
Ajax Reef	0.050 (0.010)	0	0	0
North of Ajax Reef	0	0	0	0
Star Reef	0	0	0	0
Ledberry Reef	0	0	0	0
East of Biscayne Bay	0.017 (0.001)	0	0	0
<i>Low-relief spur and groove</i>				
Offshore Pacific Reef	0	0	0	0
Between Star & Triumph Reef	0.033 (0.001)	0	0	0
SW of Brewster Reef	0	0	0	0.017 (0.001)
South of Fowey Rocks	0.033 (0.004)	0	0	0
North Dixie Shoal	0.033 (0.001)	0.050 (0.004)	0	0
French Reef **	0	0	0	0
East of Pelican Shoal	0	0	0.017 (0.001)	0
Eastern Sambo**	0	0	0	0
Western Sambo*	0	0	0	0
Between W. Sambo & E. Dry Rocks	0.017 (0.001)	0	0	0

Table 12. Mean (1 SE) densities (no. individuals per m²) of incidental marine invertebrates from surveys of four 15 m² plots per survey location. Noted are sites within Sanctuary no-take zones (*) and sites adjacent and seaward of the zones (**). Listed are only those sites surveyed where species were found within four 15 m² plots at each site.

Habitat type/site location	<i>Astrophyton muricatum</i>	<i>Periclemenes pedersoni</i>	<i>Stenopus hispidus</i>	<i>Stenorhyncus seticornis</i>
<i>Offshore patch reef</i>				
Inshore of Pacific Reef	0	0	0.033 (0.002)	0
<i>Patchy hard-bottom in sand</i>				
Between Star & Triumph Reef	0.133 (0.009)	0	0	0.017 (0.001)
North of Fowey Rocks	0.233 (0.016)	0	0	0
<i>Low-relief hard bottom</i>				
Ajax Reef	0	0	0	0
North of Ajax Reef	0.067 (0.003)	0	0	0
Star Reef	0.133 (0.033)	0	0	0
Ledberry Reef	0.333 (0.086)	0	0	0.017 (0.001)
East of Biscayne Bay	0	0	0.017 (0.001)	0.033 (0.004)
<i>Low-relief spur and groove</i>				
Offshore Pacific Reef	0.067 (0.018)	0	0	0.017 (0.001)
Between Star & Triumph Reef	0	0	0	0
SW of Brewster Reef	0	0	0	0.017 (0.001)
South of Fowey Rocks	0.083 (0.010)	0.033 (0.004)	0.017 (0.001)	0.033 (0.002)
North Dixie Shoal	0.083 (0.010)	0.017 (0.001)	0.017 (0.001)	0.033 (0.002)
French Reef**	0	0	0	0.017 (0.001)
East of Pelican Shoal	0.017 (0.001)	0	0	0
Eastern Sambo**	0.017 (0.001)	0	0	0
Western Sambo*	0	0	0	0.017 (0.001)
Between W. Sambo & E. Dry Rocks	0	0	0	0