

Phorbasterones A–D, Cytotoxic *Nor*-Ring A Steroids from the Sponge *Phorbas amaranthus*

Makoto N. Masuno,[†] Joseph R. Pawlik,[‡] and Tadeusz F. Molinski*[†]

Department of Chemistry, University of California, Davis, California 95616, and Biological Sciences and Center for Marine Science Research, University of North Carolina at Wilmington, North Carolina 28403-3297

Received October 17, 2003

The sponge *Phorbas amaranthus* from Florida contains the new ring A-contracted steroids, phorbasterones A–D, and the known anthosterones A and B. The structures of phorbasterones A–D were determined by interpretation of their spectroscopic data. Phorbasterones show moderate cytotoxicity against HCT-116 tumor cells.

In a systematic survey of Caribbean sponges, Pawlik and co-workers found that extracts of the bright-red sponge *Phorbas amaranthus* deterred feeding by the bluehead wrasse *Thalassoma bifasciatum*.¹ The measured nutrient content of *P. amaranthus* was significant, but the tissue was exceptionally fragile (tensile strength) with no obvious physical defenses.² Therefore, the deterrent properties of the sponge are likely attributed to an undescribed “chemical defense”. In our search for the antifeedant principles of *P. amaranthus* we isolated the known oxidized steroids anthosterones A (**1**) and B (**2**) and four new congeners, phorbasterones A–D (**3–6**). Steroids **1** and **2**, with contracted cyclopentane A-rings, were first described by Anderson, Clardy, and co-workers in 1988.³ Phorbasterones comprise a family of homologues that differ from **1** and **2** by side-chain (C20–29) alkyl branching, isomerism, or oxidation level. In this report, we describe the isolation and structure elucidation of **3–6**. Metabolites from the genus *Phorbas* are rare. The only other natural products described from this genus are the alkaloids phorbazoles from a Red Sea species (*Phorbas* aff. *clathrata*),⁴ the potent cytotoxic phorboxazoles A and B from a Western Australian *Phorbas* species,^{5,6} the monocyclic diterpenoids phorbazines A and B from an Australian species,⁷ and the gagunins, which are highly oxygenated verrucosane diterpenes.⁸ Phorbasterones **3–6** were found to be cytotoxic to HCT-116 cells (IC₅₀ 1–3 μg/mL).

Samples of *P. amaranthus* collected by hand (scuba) at Dry Rock, Key Largo, Florida, were immediately frozen and kept at –20 °C until needed. The CHCl₃-soluble fraction, obtained after preliminary methanol extraction of the sponge, was purified by sequential column chromatography (silica, EtOAc–CH₂Cl₂, MeOH–CH₂Cl₂ gradient), Sephadex LH20 chromatography (hexane–CH₂Cl₂, 1:3), and reversed-phase HPLC (C₈, Microsorb, MeOH–H₂O followed by C₁₈, Microsorb, CH₃CN–H₂O) to afford anthosterones A and B (**1** and **2**) and phorbasterones A–D (**3–6**). Compounds **1** and **2** were identified by comparison of ¹H and ¹³C NMR data with those of reported values,³ while the structures of **3–6** were derived as follows.

Anthosterones **1** and **2** and the phorbasterones share a characteristic ring A-contracted steroid nucleus of general structure **7** (Figure 1). The core ¹³C NMR resonances (C1 to C19) were virtually identical for each compound (Table




Figure 1.

1). Key ¹H NMR signals, including the isolated geminal signals H₂-1 (δ 2.08, d, *J* = 13.3 Hz, 1H; 2.16, d, *J* = 13.3 Hz, 1H)³ and H-6 (δ 6.72, t, *J* = 3.3 Hz, 1H), assigned to a highly polarized α,β-unsaturated cyclopentenone and the C18 and C19 angular methyl groups (δ 0.73, s, 3H; 1.21, s, 3H), confirmed the presence of an exocyclic double bond conjugated to the cyclopentanone ring. The structural differences between **1**, **2**, and the new compounds **3–6** could be attributed solely to differences in the side chains and assigned by examination of the ¹H and ¹³C NMR spectra.

Compound **3**, C₂₉H₄₄O₄, was isomeric with **2** by a desorption electron impact mass spectrum (DEI, *m/z* 456.3247 [M⁺]). The ¹H NMR spectrum of **3** (CDCl₃) revealed signals due to a 1,2-disubstituted vinyl group (δ 5.15, m, 2H, H-22, H-23) and four methyl groups (δ 1.00, d, *J* = 6.6 Hz, 3H; 0.89, d, *J* = 6.9 Hz, 3H; 0.82, d, *J* = 6.9 Hz, 3H; 0.80, d, *J* = 6.6 Hz, 3H). Comparison of the ¹³C NMR signals of **3** C20–C28 showed an almost perfect match for (*E*)-22-

* To whom correspondence should be addressed. Tel: 530 752 6358. Fax: 530 752 8995. E-mail: tfmolinski@ucdavis.edu.

[†] University of California, Davis.

[‡] University of North Carolina at Wilmington.

Table 1. Selected ^{13}C NMR Chemical Shifts of **2**–**6** (δ , CDCl_3 , 100 MHz)

#	2 ^a	3 ^b	4 ^c	5a ^{d,h}	5b ^{e,h}	6a ^{g,h}	6b ^{f,h}
2	79.9	79.9	79.9	79.9	79.9	79.9	79.9
3	173.4	173.4	173.4	173.4	173.4	173.4	173.4
4	200.9	200.9	200.9	200.9	200.9	200.9	200.9
5	145.1	145.1	145.1	145.1	145.1	145.1	145.1
6	136.3	136.3	136.3	136.3	136.3	136.3	136.3
20	35.7	40.2	35.7	40.4	40.4	36.1	36.2
		(-0.1)	(0.0)	(-0.1)	(0.0)	(0.0)	(0.0)
21	21.9	21.0	18.7	21.1	20.9	18.8	18.8
		(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)
22	31.0	135.8	36.1	138.1	138.0	33.9	33.9
		(-0.2)	(0.0)	(-0.3)	(-0.2)	(0.0)	(0.0)
23	34.6	132.1	23.8	129.5	129.5	26.1	26.4
		(+0.2)	(0.0)	(+0.2)	(+0.2)	(0.0)	(0.0)
24	156.8	43.1	39.5	51.2	51.2	45.8	46.0
		(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)
25	33.8	33.2	28.0	31.9	31.8	29.1	28.9
		(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)
26	21.9	19.6 ⁱ	22.5 ⁱ	19.0 ⁱ	18.9 ⁱ	19.8	19.6
		(0.0)	(0.0)	(0.0)	(0.0)	(-0.1)	(0.0)
27	21.8	20.2 ⁱ	22.8 ⁱ	21.3 ⁱ	21.2 ⁱ	19.0 ⁱ	19.0 ⁱ
		(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)
28	106.0	18.0		25.4 ⁱ	25.4 ⁱ	23.0 ⁱ	23.0 ⁱ
		(0.0)		(0.0)	(0.0)	(-0.1)	(0.0)
29				12.2	12.4	12.3	12.3
				(0.0)	(0.0)	(0.0)	(0.0)
MeO	53.4	53.4	53.4	53.4	53.4	53.4	53.4

^a The numbering for anthosterone B (ref 1) is changed here to be consistent with that for **3**–**6**. Differences in δ ($\Delta\delta$) for side-chain resonances (in parentheses) were computed with respect to the following reference compounds. ^b 22-Dehydrocampesterol. ^c 5 α -Cholestan-3-one. ^d Stigmasterol. ^e Poriferasterol. ^f Clionasterol. ^g Sitosterol as follows: [$\delta_{\text{phorbasterone}} - \delta_{\text{standard sterol}}$]. ^h Isolated as a 1:1 mixture. ⁱ Interchangeable.

dehydrocampesterol,^{9,10} which confirms the double-bond location (C22–C23) and configuration in **3**.

Phorbasterone B (**4**), $\text{C}_{28}\text{H}_{44}\text{O}_4$, is a lower homologue of **3** (DEI, m/z 444.3240 [M^+]) that contains one less double bond than **1**, as evidenced by lack of side-chain vinyl signals in the ^1H NMR spectrum of **4** (CDCl_3). Comparison of side-chain ^{13}C NMR signals, particularly of C20–C27, and the ^1H NMR methyl signals (δ 0.93, d, $J = 6.6$ Hz, 3H; 0.87, d, $J = 6.6$ Hz, 3H; 0.86, d, $J = 6.6$ Hz, 3H) matched the signals of a 5 α -cholestan-3-one.^{10,11} Thus, phorbasterone B (**4**) is 22,23-dihydroanthosterone A.

Using reversed-phase HPLC under varying conditions we effected a separation of phorbasterones C (**5**) and D (**6**), each as a mixture of epimers. Accurate mass measurement of **5** (DEI m/z 470.3382 [M^+]) revealed a formula $\text{C}_{30}\text{H}_{46}\text{O}_4$ that corresponds to an ethyl-branched homologue of **3**, while the formula of **6**, $\text{C}_{30}\text{H}_{48}\text{O}_4$ (DEI, m/z 472.3552 [M^+]), is the 22,23-dihydro derivative of **5**. These structural differences between **5** and **6** were fully supported by an analysis similar to that described for **3** and **4**; however, an observed doubling of side-chain signals in the ^{13}C NMR spectra suggested that **5** and **6** were each an inseparable 1:1 epimeric mixture, most likely at C-24, from biosynthetic precedents (*cf.* sitosterol and clionasterol, *vide infra*). Consequently, phorbasterones C and D were each characterized as a 1:1 epimeric mixture at C-24 (epimers are indicated by suffixes **a** and **b**).

Analysis of the ^1H NMR spectrum of **5** (CDCl_3) supported a disubstituted *E*-double bond at C22,23 (δ 5.14, ddd, $J = 2.8, 8.4, 15.2$ Hz, 1H; 5.01, ddd, $J = 2.0, 8.4, 15.2$ Hz, 1H). Inspection of the DQF-COSY spectrum showed four methyl groups (δ 1.02, d, $J = 6.4$ Hz, 3H; 0.84–0.76, overlapped, m, 9H), one of which was part of an ethyl group, as suggested by a cross-peak between an overlapped CH_3 signal and a CH_2 signal at δ 1.65 (m). However, the ^1H

NMR signals were not sufficiently dispersed to allow accurate ^1H NMR chemical shift or coupling constant measurements. Instead, examination of the ^{13}C NMR spectra of **5**, including the DEPT spectra, revealed the expected additional CH_2 (δ 25.4, t, C28) and exceptionally high-field CH_3 signal (δ 12.2, q, C29) of the ethyl group and allowed complete assignment of the structures of **5** and **6**, including stereochemistry, as follows.

The stereochemical assignment of the C-24 configuration followed from careful pairwise comparison and least-difference analysis of ^{13}C NMR chemical shifts in **5a** and **5b** with those of the assigned side-chain signals of the known C-24 epimers, stigmasterol^{10,12} and poriferasterol^{10,13} (Table 1). Since *diastereomeric* differences observed in the side-chain ^{13}C NMR signals are most likely influenced by the nearest stereogenic center, C-20 (which is invariably 20*R* in natural sterols), the pairwise analysis of ^{13}C NMR chemical shifts allows assignment of *absolute* C-24 stereochemistry in **5a** and **5b** as 24*S* and 24*R*, respectively. Similarly, pairwise comparison of ^{13}C NMR signals of **6** with those of sitosterol^{10,14} and its C-24 epimer clionasterol^{10,15} allowed assignment of the 24*R* and 24*S* configurations to the epimers of phorbasterone D, **6a** and **6b**, respectively (note the change in CIP priorities at C24).

Phorbasterones A–D (**3**–**6**) showed moderate cytotoxicity (IC_{50} 1–3 $\mu\text{g}/\text{mL}$) toward cultured HCT-116 colon tumor cells. The solvent fraction from which compounds **1**–**6** were derived was not active as a fish feeding deterrent in assays with *Thalassoma bifasciatum*.

In summary, we have identified four new ring A-contracted steroids, phorbasterones A–D (**3**–**6**), from *P. amaranthus*. Work on the chemical nature of the feeding deterrent principles is ongoing and will be reported in due course.

Experimental Section

Experimental Procedures. General procedures are described elsewhere.¹⁶ Mass spectrometric measurements were performed at University of California, Riverside Mass Spectrometry Facility. NMR measurements were carried out on a Varian Inova 400 MHz NMR spectrometer equipped with either a $\{^1\text{H}\{^{15}\text{N}-^{31}\text{P}\}$ pulsed-field gradient (PFG) indirect-detection probe or $^1\text{H}/^{13}\text{C}/^{15}\text{N}/^{31}\text{P}$ PFG auto-switchable probe. DQF-COSY experiments were carried out with gradient-enhanced pulse sequences.

Animal Material. *Phorbis amaranthus* (02-13-054) was collected by hand using scuba at -3 to -10 m at North Dry Rocks, Key Largo, FL (25°07.850' N, 080°17.521' W) and identified by one of the authors, J.R.P. The sponge was stored for 2 months at -20 °C before extraction. A voucher specimen stored at UNC Wilmington is available from J.R.P.

Collection and Extraction of *P. amaranthus*. The lyophilized tissue (339 g) was gently agitated in MeOH (800 mL) and H_2O (200 mL) using an overhead stirrer (5 °C for 24 h). After filtration, extraction of the tissue was repeated twice with fresh MeOH (900 mL) and H_2O (100 mL), and a third time with MeOH (100 mL) and CHCl_3 (900 mL). Removal of volatiles from the CHCl_3 -MeOH extract gave a deep-red gum (9.3 g). The majority of the extract (9.0 g) was applied to a silica column and eluted with a gradient (EtOAc in CHCl_3 , then MeOH in CHCl_3). The 20% EtOAc fraction contained the crude anthosterones **1** and **2** and phorbasterones **3**–**6**. This fraction was further separated on Sephadex LH-20 (1:3 hexane- CH_2Cl_2) to yield a purified fraction of **1**–**6** (90 mg). Final purification was achieved by reversed-phase HPLC (C_8 Microsorb 90:10 MeOH- H_2O , then C_{18} , Microsorb, $\text{CH}_3\text{CN}-\text{H}_2\text{O}$) to afford, in order of elution, anthosterone A (**1**, 1.5 mg, 0.00037% dry wt), anthosterone B (**2**, 2.4 mg, 0.00059% dry wt), phorbasterone A (**3**, 4.4 mg, 0.0011% dry wt), phorbasterone

Table 2. Selected ¹H NMR Chemical Shifts of **2–6** (δ, CDCl₃)^a

#	2	3	4	5^d	6
1	2.10, d, <i>J</i> = 13.5 Hz 2.18, d, <i>J</i> = 13.5 Hz	2.08, d, <i>J</i> = 13.8 Hz 2.17, d, <i>J</i> = 13.8 Hz	2.09, d, <i>J</i> = 13.8 Hz 2.18, d, <i>J</i> = 13.8 Hz	2.08, d, <i>J</i> = 14 Hz 2.17, d, <i>J</i> = 14 Hz	2.08, d, <i>J</i> = 13.6 Hz 2.17, d, <i>J</i> = 13.6 Hz
6	6.73, t, <i>J</i> = 3.3 Hz	6.71, t, <i>J</i> = 3.6 Hz	6.73, t, <i>J</i> = 3.3 Hz	6.72, t, <i>J</i> = 3.6 Hz	6.72, t, <i>J</i> = 3.6 Hz
7	2.38, ddd, <i>J</i> = 4.0, 6.0, 21.0 <i>c</i>	2.38, ddd, <i>J</i> = 3.9, 6.0, 20.7 <i>c</i>	2.40, ddd, <i>J</i> = 3.9, 6.3, 20.7 <i>c</i>	2.37, ddd, <i>J</i> = 3.6, 5.7, 21.0 1.85, ddd, <i>J</i> = 3.6, 9.3, 21.0	2.38, ddd, <i>J</i> = 4.0, 6.0, 21.0 1.85, ddd, <i>J</i> = 4.0, 9.0, 21.0
18	0.73, s	0.72, s	0.72, s	0.72, s	0.71, s
19	1.20, s	1.20, s	1.20, s	1.20, s	1.18, s
21	0.96, d, <i>J</i> = 6.7 Hz	1.00, d, <i>J</i> = 6.6 Hz	0.93, d, <i>J</i> = 6.6 Hz	1.02, d, <i>J</i> = 6.4 Hz	0.91, d, <i>J</i> = 6.4 Hz
22	<i>c</i>	5.13–5.17, m	<i>c</i>	5.14, ddd, <i>J</i> = 2.8, 8.4, 15.2 Hz	<i>c</i>
23	<i>c</i>	5.13–5.17, m	<i>c</i>	5.01, ddd, <i>J</i> = 2.8, 8.4, 15.2 Hz	<i>c</i>
26 ^b	1.04, d, <i>J</i> = 6.5 Hz	0.89, d, <i>J</i> = 6.9 Hz	0.87, d, <i>J</i> = 6.6 Hz	0.83, d, <i>J</i> = 6.6 Hz	0.82, d, <i>J</i> = 6.6 Hz
27 ^b	1.04, d, <i>J</i> = 6.5 Hz	0.82, d, <i>J</i> = 6.9 Hz	0.86, d, <i>J</i> = 6.6 Hz	0.79, d, <i>J</i> = 6.6 Hz	0.79, d, <i>J</i> = 6.6 Hz
28		0.80, d, <i>J</i> = 6.6 Hz	-	<i>c</i>	<i>c</i>
OH	3.78, brd, <i>J</i> = 1.2 Hz	3.78, brd, <i>J</i> = 0.9 Hz	3.80, brs	3.78, brd, <i>J</i> = 0.9 Hz	3.78, brs
MeO	3.77, s	3.76, s	3.78	3.76	3.76

^a Spectra of **2** and **6** were recorded at 400 MHz, while compounds **3**, **5**, and **6** were recorded at 300 MHz. ^b Chemical shifts for H26 and H27 are interchangeable. ^c Unresolved. ^d Assignments from DQFCOSY (600 MHz).

terone B (**4**, 6.7 mg, 0.0017% dry wt), phorbasterone C (**5**, 1.1 mg, 0.0003% dry wt), and phorbasterone D (**6**, 0.7 mg, 0.0002% dry wt).

(–)-**Phorbasterone A (3)**: colorless solid, [α]_D –45.4° (*c* 0.19, CHCl₃); UV (CH₃CN) λ_{max} 250 nm (ε 9600); IR (film) ν_{max} 3453, 2956, 1745, 1720, 1651 cm⁻¹; ¹H NMR (CDCl₃, see Table 2); ¹³C NMR (100 MHz, CDCl₃, see Table 1); HRMS (DEI) *m/z* 456.3247 [M⁺] (calcd for C₂₉H₄₄O₄, 456.3239).

(–)-**Phorbasterone B (4)**: colorless solid, [α]_D –54.6° (*c* 0.28, CHCl₃); UV (CH₃CN) λ_{max} 250 nm (ε 9300); IR (film) ν_{max} 3469, 2952, 1745, 1720, 1650 cm⁻¹; ¹H NMR (CDCl₃, see Table 2); ¹³C NMR (100 MHz, CDCl₃, see Table 1); HRMS (DEI) *m/z* 444.3240 [M⁺] (calcd for C₂₈H₄₄O₄, 444.3239).

Phorbasterone C (5a and 5b): colorless solid, UV (CH₃CN) λ_{max} 250 nm (ε 9100); IR (film) ν_{max} 3357, 2958, 1745, 1720, 1650, 1384 cm⁻¹; ¹H NMR (CDCl₃, see Table 2); ¹³C NMR (100 MHz, CDCl₃, see Table 1); HRMS (DEI) *m/z* 470.3382 [M⁺] (calcd for C₃₀H₄₆O₄, 470.3396).

Phorbasterone D (6a and 6b): colorless solid, UV (CH₃CN) λ_{max} 250 nm (ε 9000); IR (film) ν_{max} 3357, 2958, 1745, 1720, 1650 cm⁻¹; ¹H NMR (CDCl₃, see Table 2); ¹³C NMR (100 MHz, CDCl₃, see Table 1); HRMS (DEI) *m/z* 472.3537 [M⁺] (calcd for C₃₀H₄₈O₄, 472.3552).

Cytotoxicity Assays. Cytotoxicity was measured with HCT-116 cells using the MTS method.¹⁷ Briefly, compounds were assayed with compounds in DMSO (final concentration, 1% v/v) and run against etoposide as positive control. HCT-116 cells were incubated in 96-well plates for 72 h before addition of MTS [(3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulphophenyl)-2H-tetrazolium, inner salt)]. The Promega CellTiter 96 Aqueous cell proliferation assay (Technical Bulletin No. 169) was used. Well absorbances (λ 490 nm) were corrected for background and expressed as a percentage of the negative control (DMSO, only).

Acknowledgment. We thank J. Cowart (University of North Carolina, Wilmington) for assistance with collection of

Phorbasp. sp., and S. Kelly (Scripps Institution of Oceanography) and S. Lievens (UCD) for performing the HCT-116 bioassays. The Bruker Avance 600 MHz spectrometer was purchased with funds provided by NIH RR11973. The Varian Inova 400 MHz spectrometer was purchased with funds provided by NSF CHE-9808183. This work was supported, in part, by grants from NIH (CA 85602) to T.F.M., and the NSF Biological Oceanography Program (OCE-0095724) and NOAA/National Undersea Research Center at UNCW to J.R.P.

References and Notes

- Pawlik, J. R.; Chanas, B.; Toonen, R. J.; Fenical, W. *Mar. Ecol. Prog. Ser.* **1995**, *127*, 183–194.
- Chanas, B.; Pawlik, J. R. *Mar. Ecol. Prog. Ser.* **1995**, *127*, 195–211.
- Tischler, M.; Ayer, S. W.; Anderson, R. J.; Mitchell, J. F.; Clardy, J. *Can. J. Chem.* **1988**, *66*, 1173–1178.
- Rudi, A.; Stein, Z.; Green, S.; Goldberg, I.; Kashman, Y.; Benayahu, Y.; Schleyer, M. *Tetrahedron Lett.* **1994**, *35*, 2589–2592.
- Searle, P. A.; Molinski, T. F. *J. Am. Chem. Soc.* **1995**, *117*, 8126–8131.
- Searle, P. A.; Molinski, T. F.; Brzezinski, L. J.; Leahy, J. W. *J. Am. Chem. Soc.* **1996**, *118*, 9422–9423.
- McNally, M.; Capon, R. *J. Aust. J. Chem.* **2001**, *64*, 645–647.
- Rho, J.-R.; Lee, H.-S.; Sim, C. J.; Shin, J. *Tetrahedron* **2002**, *58*, 9585–9591.
- Findlay, J. A.; Patil, A. D. *Can. J. Chem.* **1985**, *63*, 2406–2410.
- Wright, J. L. C.; McInnes, A. G.; Shimizu, S.; Smith, D. G.; Walter, J. A.; Idler, D.; Khalil, W. *Can. J. Chem.* **1978**, *56*, 1898–1903.
- Blunt, J. W.; Stothers, J. B. *Org. Magn. Reson.* **1977**, *9*, 439–464.
- Holland, H. L.; Diakow, P. R. P.; Taylor, G. L. *Can. J. Chem.* **1978**, *56*, 3121–3127.
- Nicotra, F.; Ronchetti, F.; Russo, G.; Toma, L.; Ranzi, B. M. *Magn. Reson. Chem.* **1985**, *23*, 134–136.
- Nes, W. D.; Norton, R. A.; Benson, M. *Phytochemistry* **1992**, *31*, 805–811.
- Reynolds, W. F.; McLean, S.; Tay, L.-L.; Yu, M.; Enriquez, R. G.; Estwick, D. M.; Pascoe, K. O. *Magn. Reson. Chem.* **1997**, *35*, 455–462.
- Searle, P. A.; Richter, R. K.; Molinski, T. F. *J. Org. Chem.* **1996**, *61*, 4073–4079.
- Zhou, G.-X.; Molinski, T. F. *Mar. Drugs* **2003**, *1* (1), 46–53.

NP034037J