



The American Midland Naturalist

An International Journal of
Ecology, Evolution and Environment

Effects of White-Tailed Deer (*Odocoileus virginianus*) on the Maritime Forest of Bald Head Island, North Carolina

JOHN TAGGART¹ AND ZACHARY LONG
University of North Carolina Wilmington 28403



Effects of White-Tailed Deer (*Odocoileus virginianus*) on the Maritime Forest of Bald Head Island, North Carolina

JOHN TAGGART¹ AND ZACHARY LONG

University of North Carolina Wilmington 28403

ABSTRACT.—Overabundance and associated impacts of deer on mainland forests of the United States have been topics of extensive investigation and management discussion. Conversely, deer populations on barrier islands have been less studied with few investigations of their impacts on maritime evergreen forest. We investigated how deer influenced understory vegetation and oak seedling survival following culling of deer. In 2007 twenty 10 m × 10 m paired exclosure and control plots were established within a 77.3 ha protected area on Bald Head Island, North Carolina. Vegetation was sampled in 2011. We determined deer did not affect understory vegetation and oak survival was low and not influenced by deer. Chronic over-browsing (*i.e.*, ‘ghost of herbivory past’) was unlikely because deer were not abundant on Bald Head Island until the 1990s, oak seedlings were observed where light was abundant, and deer were observed using other cover types on the island. Continued monitoring of the forest understory and maintenance of deer at the current population level will help conserve maritime forest on Bald Head Island.

INTRODUCTION

Throughout the eastern United States, the abundance of white-tailed deer (*Odocoileus virginianus*) has increased dramatically over the last century (McShea *et al.*, 1997; Russel *et al.*, 2001). This could have occurred because of increased habitat fragmentation from land development or reduced predators and hunting pressure (Augustine and DeCalesta, 2003; Knight *et al.*, 2009). Regardless of the cause, high deer densities can have significant effects on plant communities (Côté *et al.*, 2004) and have changed floristic compositions by over-browsing of understory vegetation or selective browsing of species that inhibited forest succession (Augustine and Frelich, 1998; Russell *et al.*, 2001; Horsely *et al.*, 2003; Côté *et al.*, 2004; Long *et al.*, 2007; Baiser *et al.*, 2008; Knight *et al.*, 2009; Aronson and Handel, 2011).

Plant communities on southeastern U.S.A. barrier islands have been particularly susceptible to browsing because they grow in nutrient-poor soils and are subjected to additional physical stress such as salt spray and shifting sediments (Au, 1974; Ray *et al.*, 2001). Maritime forest regeneration and maintenance of community structure has occurred in response to the physical environment and disturbance regime (Au, 1974; Turner and Bratton, 1987; Conner *et al.*, 2005). Despite increases in deer abundances on southeastern barrier islands (Osborne *et al.*, 1992; Ray *et al.*, 2001), few studies investigated how herbivores influenced maritime forest dynamics (Turner and Bratton, 1987; Forrester *et al.*, 2006).

Deer presence on Smith Island in North Carolina (Fig. 1) has been relatively recent as deer were not recorded in mammalian surveys conducted in 1964 (Cooper and Satterthwaite, 1964) or 1970 (Parnell and Adams, 1970). Deer colonized the island in the mid-1980s and established a viable population in the early 1990s (Ray *et al.*, 2001). Studies have suggested barrier islands can sustain deer populations ≤ 19 deer/km² (Stransky, 1969;

¹ Corresponding author present address: Department of Environmental Studies, University of North Carolina Wilmington 28403. Telephone: (910) 962-7909; e-mail: taggartj@uncw.edu



FIG. 1.—Location of Bald Head Island and the maritime forest reserve within the Smith Island complex that includes Middle and Bluff islands

Osborne *et al.*, 1992; Sherrill *et al.*, 2010). Since 1999 surveys showed deer densities on the Bald Head Island portion of Smith Island increased dramatically, fluctuated considerably, and exceeded 19 deer/km² most years (Sherrill *et al.*, 2010; Bald Head Island Conservancy, 2012; Table 1). Complaints of browsing damage to residential plantings and several deer deaths from vehicular accidents (*e.g.*, construction trucks) during 1999–2002 prompted the local government, Bald Head Island Village Council (2014), to hire sharpshooters for culling (Bald Head Island Conservancy, 2012). Culls were conducted in 2002, 2005, 2006, 2008, and 2011, but the village council was concerned about deer impacts on the local forest preserve. Our objective was to determine if deer influenced dynamics of a protected maritime evergreen forest – south Atlantic subtype (Schafale, 2012) on a barrier island in the southeastern U.S.A.

TABLE 1.—Deer Population Estimates for Bald Head Island 1999–2011 (Sherrill *et al.*, 2010; Bald Head Island Conservancy, 2012). Estimates from Sherrill *et al.*, 2010 are listed first for 2008 and 2009; others are from Bald Head Island 2012

Year	Number of deer	Number culled	Density (km ²)
1999	93		15.0
2000	185		29.8
2001	316		51.0
2002	257*	149	41.5
2003	54		8.7
2004	332		53.5
2005	359*	100	57.9
2006	256*	145	41.3
2007	134		21.6
2008	107/148*	71	17/23.9
2009	93/97		15/15.6
2010	141		22.8
2011	155*	73	25.0
2012	76		12.3

* - herd reduction performed

STUDY AREA

The study area was within the Bald Head Woods Coastal Reserve (33°51'10.80"N, 77°58'30.68"W), a 77.3 ha maritime forest preserve located on Bald Head Island in North Carolina (Fig. 1). Bald Head Island consists of approximately 620 ha of ocean beach to sound communities on a barrier island where the majority of uplands are occupied by privately-owned tracts with special attention paid to protection of natural features. This area represents the southern portion of Smith Island, a complex which also encompasses Middle Island, Bluff Island, and other beaches/marshes in private and public ownerships.

METHODS

In 2007 we established paired 20 plots that each consisted of a 10 m × 10 m enclosure and a 10 m × 10 m control. This plot size and square configuration has been shown to provide sufficiently detailed information on terrestrial vegetation composition and structure for diverse applications and compatibility with data derived from commonly used methods (Peet *et al.*, 1998). We selected sites for paired plots to represent variation in species composition, physical structure of forest regeneration phases, and environmental gradients that trended from southwest to northeast (*i.e.*, a gradient direction that included younger to older dunes, increasing moisture availability, and decreasing salt spray) (Brewer, 2008). Each enclosure was surrounded by a black polypropylene mesh (4.4 cm × 4.4 cm) deer fence 2.13 m in height and 12 m × 12 m in area to avoid edge effect on the interior 100 m² sample. At least one gap (10 cm high × 25 cm long) was created along the bottom of each side to allow small animal access. This design was consistent with deer enclosures of this size (Nicholas *et al.*, 2007; Deer Facts, 2009) and successfully prevented deer access.

In 2011 we sampled each plot to acquire tree, shrub, and vine data. We used Weakley (2012) as the plant identification guide and deposited a voucher specimen in the University of North Carolina Wilmington Herbarium (WNC) for each vascular plant species sampled. Trees, shrubs, and vines were divided into three height classes: small (<50 cm), medium (51–130 cm), and large (>131 cm). We measured diameter at breast height of all other

species >1.3 m. Herb occurrences in exclosures and throughout the maritime forest were sparse and not included. The lack of herbaceous vegetation in North Carolina maritime forests was reported by Bordeau and Oosting (1959) and Au (1974). Paucity of herbs in this forest was likely due to low light conditions under the closed canopy. In a vegetation survey conducted on this island prior to deer colonization, Bordeau and Oosting (1959) reported: "...the herb layer was sparse, except below openings in the canopy."

We used paired *t*-tests to determine overall density of woody species, density of the most abundant woody understory species, and understory diversity of woody species (Shannon index). We examined the influence of deer on understory composition by permutated analysis of variance (PERMANOVA). We conducted all analyses in R version 2.15.0 (R Development Core Team, 2012).

We assessed deer impacts on young live oaks (*Quercus virginiana*) by transplanting seedlings and saplings derived from acorns harvested on Bald Head Island. In 2010 we evenly divided and planted 144 seedlings and 72 saplings (2 to 3 y old) within six exclosures and six control plots. In 2011 we evenly divided another 110 seedlings between an exclosure and a single control. We identified initial transplants by individual tags (*i.e.*, coded 'twist ties') and small survey flags for groups of three seedlings. We planted all other seedlings in rows identified by numbered plant stakes. Twelve months following transplantation, we attempted to locate seedlings and saplings for assessment of survival.

To sample ages of canopy recruitment, we obtained tree cores from dominant oaks in all plots (0.4 ha total area). Live oaks were too large in diameter and contained decomposed heartwood inadequate for age determination, but we sampled all sand laurel oaks (*Q. hemisphaerica*). We sanded and examined each core with a dissecting microscope to determine tree age.

RESULTS

Understory tree, shrub, and vine species densities were similar between exclosure and control plots in the small [$t(19) = 0.64$, $P = 0.53$], medium [$t(19) = 0.14$, $P = 0.89$], and large size classes [$t(19) = 0.52$, $P = 0.60$] after 4 y of exclusion (Fig. 2). The understory was dominated by yaupon (*Ilex vomitoria*) with lesser amounts of red bay (*Persea borbonia*) and cabbage palmetto (*Sabal palmetto*) (Table 2). Yaupon density was similar in all size classes [$t(19) = 0.92$, $P = 0.37$ for small, $t(19) = 0.88$, $P = 0.39$ for medium, $t(19) = 0.81$, $P = 0.42$ for large]. Shannon diversity was similar between exclosures and controls in all size classes of understory species [$t(19) = -1.24$, $P = 0.23$ for small, $t(19) = -1.78$, $P = 0.09$ for medium, $t(19) = -1.55$, $P = 0.14$ for large, Fig. 3]. Species composition was similar for small (F Model = 0.43, $r^2 = 0.01$, $P = 0.78$), medium (F Model = 0.62, $r^2 = 0.02$, $P = 0.60$), and large size classes (F Model = 0.28, $r^2 = 0.01$, $P = 0.81$, Fig. 4).

The canopy consisted of oaks with only sporadic occurrences of American holly (*Ilex opaca* var. *opaca*), southern red cedar (*Juniperus virginiana* var. *silicicola*), and Carolina laurel cherry (*Prunus caroliniana*) (Table 3). Although cabbage palmetto was common in the understory, it was absent from the canopy within plots.

Only six of the 127 transplanted seedlings survived in the control plots and none of the 127 survived in the exclosures. Three of the 36 transplanted saplings survived in the controls, while zero survived in the exclosures. Evidence of approximately 30 plants removed or killed by digging, presumably by small animals, was observed during monitoring. Of 17 sand laurel oak canopy and subcanopy trees cored, ring counts indicated a mean of 46 y (SE = 3.38) and tree age ranged from 23 to 73 y.




FIG. 2.—Stem density in the different size classes and total stem density in exclosures and controls at Bald Head Island, 2011. The box is bordered on the lower end by 25% quartile and on the upper end by the 75% quartile. The solid line represents the median diversity. The lower and upper lines represent the minimum and maximum values, respectively

DISCUSSION

Overabundance and associated impacts of deer on mainland forests of the United States have been topics of extensive investigation and management discussion (Harlow *et al.*, 1970; Stromayer and Warren, 1997; Russell *et al.*, 2001; Rooney and Waller, 2003; Côté *et al.*, 2004; Webster *et al.*, 2005; Tanentzap *et al.*, 2011). Deer populations on barrier islands, however, have been considerably less studied (Stransky, 1969; Epstein *et al.*, 1983, 1985; Miller, 1988; Rowland, 1989; Osbourne *et al.*, 1992; and Ray *et al.*, 2001), with few studies concerning impacts on maritime evergreen forest (*e.g.*, Bratton and Kramer, 1993). Because maritime forest area has been drastically reduced in recent decades by development and other land conversion activities, protection and stewardship of remaining tracts was a priority in southeastern coastal states (Lopazanski *et al.*, 1988; Bellis, 1995; North Carolina Coastal Reserve, 2014). Our results indicated the deer population on Bald Head Island had

TABLE 2.—Relative abundances of each species found in maritime forest understory by size class, Bald Head Island, 2011. There were 14,842 individuals in the small size class, 6246 in the medium size class, and 3293 in the large size class

Species	Small	Medium	Large	Total
<i>Berchemia scandens</i>	<0.01	<0.01	0	<0.01
<i>Carpinus caroliniana</i>	0	0	0	0
<i>Cartrema americana</i>	0	0	0	0
<i>Cornus florida</i>	0	0	0	0
<i>Gelsemium sempervirens</i>	<0.01	0	0	<0.01
<i>Ilex opaca</i> var. <i>opaca</i>	0	0	0	0
<i>Ilex vomitoria</i>	0.85	0.71	0.66	0.79
<i>Juniperus virginiana</i> var. <i>silicicola</i>	<0.01	0	0	<0.01
<i>Muscadinia rotundifolia</i>	<0.01	<0.01	0	<0.01
<i>Parthenocissus quinquefolia</i>	<0.01	0	0	<0.01
<i>Persea borbonia</i>	0.08	0.14	0.08	0.10
<i>Prunus caroliniana</i>	0	0.03	0.01	0.01
<i>Quercus hemisphaerica</i>	<0.01	0	0	<0.01
<i>Quercus virginiana</i>	<0.01	0	0	<0.01
<i>Sabal palmetto</i>	0.06	0.11	0.25	0.10
<i>Smilax bona-nox</i>	<0.01	0	<0.01	<0.01
<i>Toxicodendron radicans</i>	<0.01	0	0	<0.01




FIG. 3.—Shannon understory diversity within 20 exclosure and control plots located in the Bald Head Island maritime forest, 2011




FIG. 4.—Nonmetric multidimensional scaling analyses of small, medium, and large (left to right) understory size classes in enclosure and control plots, Bald Head Island, 2011

relatively little influence on the understory of a maritime evergreen forest. We may have been unable to detect an effect because: (1) over-browsing for many years has eliminated any potential for recovery of plant communities by eliminating the seed bank (“the ghost of herbivory past” sensu, Carson *et al.*, 2005), (2) deer may use the forest as cover, but not browse, and forest dynamics are driven by disturbance, or (3) the deer population has been reduced to levels that allow plants to persist in the understory.

DOES ‘THE GHOST OF HERBIVORY PAST’ HAUNT BALD HEAD ISLAND?

If over-browsing by deer occurs over a sufficiently long time period, highly selected browse species may be eliminated from the understory and seed bank. Over-browsing can lead to “recalcitrant understory layers” dominated by species with low selectivity at the expense of original species diversity (Côté *et al.*, 2004; Royo and Carson, 2006). “The ghost of herbivory past” suggests that in the absence of a seedling and seed bank, it may take years for locally extirpated plant species to recolonize after deer numbers are lowered or deer are excluded from an area (Carson *et al.*, 2005). However, we do not believe this phenomenon contributed to the lack of differences between exclosures and controls in stem density, diversity, and community composition in this study. Yaupon, the dominant understory species, is considered a highly selected browse species and red bay, the second most abundant understory species, is intermediate in palatability (Goodrum and Reid, 1958; Shadow, 2011; U.S. Forest Service, 2012).

Previous surveys of the Bald Head Island maritime forest (Coker, 1918; Wells, 1932; Bourdeau and Oosting, 1959) included the same woody species, sparse herb stratum, and overall characteristics that we detected in this study. Also, the understory composition and structure that we sampled were similar to quantitative data from plots sampled in 1988 by the Carolina Vegetation survey (2013) in the same protected forest. Our sample size ($n =$

TABLE 3.—Relative abundances and average DBHs of canopy species sampled in the Bald Head Island maritime forest, 2011. A total of 28 trees were sampled in all plots (0.4 ha)

Species	Relative abundance	Mean DBH (cm)	SE
<i>Ilex opaca</i> var. <i>opaca</i>	0.03	54.2	0.00
<i>Juniperus virginiana</i> var. <i>silicicola</i>	0.06	40.1	1.65
<i>Prunus caroliniana</i>	0.06	24.4	1.66
<i>Quercus hemisphaerica</i>	0.61	30.9	6.60
<i>Quercus virginiana</i>	0.24	44.9	4.36

20) was much higher than previous studies that found significant results (*e.g.*, $n = 6$ in Long *et al.*, 2007). The large sample size and paired design suggested, if deer significantly affected understory composition, we would have detected the change. We also noted the lack of herbaceous vegetation in the controls and exclosures likely indicated light limitation because a survey conducted on this island prior to deer colonization (Bourdeau and Oosting, 1959) found herbs were restricted to canopy gaps. The lack of change in browse species (*i.e.*, yaupon and red bay) following deer exclusion, the consistent composition and structure of the understory over the last century, and the recent colonization (and subsequent management) of deer indicated ‘the ghost of herbivory past’ did not haunt Bald Head Island.

DO DEER CONSUME UNDERSTORY VEGETATION IN THIS FOREST?

Our results indicated deer were not over-browsing understory vegetation. In addition to the maritime forest, Bald Head Island had many other habitats including a golf course, landscaped areas, early successional areas, secondary dunes, and freshwater wetlands. During a study of deer movement and habitat selection on Bald Head Island (Sherrill *et al.*, 2010), animals were observed using a variety of habitats throughout the island (B. Sherrill, pers. comm.), but maritime forest/shrub was used by radio-collared deer at levels greater than other available cover types.

HAVE DEER BEEN CULLED TO LEVELS THAT DO NOT IMPACT UNDERSTORY VEGETATION?

Spotlight count estimates of the Bald Head Island deer population were as high as 359 individuals ($57.9/\text{km}^2$) in 2005 but have remained below 150 ($24/\text{km}^2$) individuals since 2007 (Sherrill *et al.*, 2010; Bald Head Island Conservancy, 2012; Table 1). Lack of significant changes in maritime forest vegetation and oak recruitment indicated the forest tolerated, at least in the short term, deer densities higher than predicted by Stransky (1969) for a barrier island system (*i.e.*, $15\text{--}17/\text{km}^2$). Stransky’s estimate assumed a low quality deer habitat, while some southeastern mainland forests were considered able to support approximately 19 deer/ km^2 (Stransky, 1969; Osbourne *et al.*, 1992). However, the lack of competition with other barrier island herbivores (*e.g.*, feral horses and hogs), availability of adjacent food sources, and mechanisms of oak recruitment appeared to be important mitigating factors on Bald Head Island. A target density of $15\text{--}17$ deer/ km^2 recommended by Sherrill *et al.*, 2010 was a conservative goal to ensure overall protection of island habitats.

DEER EFFECTS ON OAK RECRUITMENT

Similar to other southeastern maritime forests and previous studies on Bald Head Island (Coker, 1918; Wells, 1939; Oosting, 1954; Bourdeau and Oosting, 1959), we determined oaks dominated the Bald Head Island maritime forest canopy. Despite canopy dominance, there was a consistent lack of oak seedlings and saplings on our plots. We counted small numbers of oak seedlings even though several plots encompassed mature trees that produced acorns during this study.

The high mortality of live oak transplants in exclosure and control plots that we observed tended to confirm results of Warbington-Wells (2004) who noted acorns planted at the soil surface in an open field had highest survivorship relative to acorns buried deeper and shaded. Even if acorns were not consumed and successfully germinated, our results indicated seedlings would not survive under a dense canopy. We observed seedling and sapling loss of 96.3% in controls and 100% in exclosures from what appeared to be digging by small animals (*e.g.*, squirrels) and shading (Bourdeau and Oosting, 1959). However, we did observe vigorous oak recruitment along island roadsides and in open areas where

sunlight and temperature were conducive for germination/growth despite the opportunities for deer to browse.

Comparisons between exclosure and control plots located within a state-managed protected area did not reveal any significant differences in understory species diversity, stem density, or composition. We believe impacts of deer on the Bald Head Island maritime evergreen forest understory during 2007–2011 were minor. Continuation of the suggested carrying capacity of 100 animals (15–17/km²) (Stransky, 1969; Sherrill *et al.*, 2010) was justified to protect the forest plus adjacent natural communities and to avoid conflicts with residential uses on the island.

Acknowledgments.—We thank staff and volunteers from the Bald Head Island Conservancy for transportation support and vital assistance with field work. Dr. Steven Brewer and his students set up all plots in 2007. The Village of Bald Head Island generously provided funding for the project. The manuscript was improved by thoughtful comments from Myla Aronson, Amy Long, Ian Thompson, and two anonymous reviewers.

LITERATURE CITED

- ARONSON, M. F. J. AND S. N. HANDEL. 2011. Deer and invasive species suppress forest herbaceous communities and canopy tree regeneration. *Nat. Areas J.*, **31**:400–407.
- AU, S.-F. 1974. Vegetation and ecological processes on Shackleford Bank, North Carolina. Issue 6, scientific monograph series. U.S. Department of the Interior, National Park Service, Washington, D.C. 86 p.
- AUGUSTINE, D. AND D. DeCALESTA. 2003. Defining deer overabundance and threats to forest communities: from individual plants to landscape structure. *Ecoscience*, **10**:472–486.
- AND L. E. FRELICH. 1998. White-tailed deer impacts on populations of an understory forb in fragmented deciduous forests. *Cons. Bio.*, **12**:995–1004.
- BAISER, B. J., L. LOCKWOOD, D. LA PUMA, AND M. F. J. ARONSON. 2008. A perfect storm: two ecosystem engineers interact to degrade deciduous forests of New Jersey. *Bio. Inv.*, **10**:785–795.
- BALD HEAD ISLAND CONSERVANCY. 2012. Bald Head Island white-tailed deer Fall 2012 population estimate. Unpublished report prepared for the Village of Bald Head Island, NC. Bald Head Island, North Carolina. 2 p. Available: <<http://www.bhic.org/media/pdf/fall%202012%20deer%20report%20dorsey.pdf>>. Accessed 9 Jul. 2014.
- BALD HEAD VILLAGE COUNCIL. 2014. Village of Bald Head Island, mission statement. Bald Head Island, NC. Available: <<http://www.villagebhi.org/government/council/index.html>>. Accessed 9 Jul. 2014.
- BELLIS, V. J. 1995. Ecology of Maritime Forests of the Southern Atlantic Coast: a community profile. Biological Report 30. U.S. Department of the Interior, National Biological Service, Washington, D.C. 107 p.
- BOURDEAU, P. F. AND H. J. OOSTING. 1959. The maritime live oak forest in North Carolina. *Ecology*, **40**:148–152.
- BRATTON, S. P. AND E. A. KRAMER. 1993. Recovery of live oak sprouts after release from browsing on Cumberland Island National Seashore, GA, p. 47–57. *In*: C. A. Cole and K. Turner (eds.). Barrier island ecology of the mid-Atlantic coast: a symposium, U.S. National Park Service Technical Report NPS/SERCAHA/NRTR – 93/04. Washington, D.C. 216 p.
- BREWER, S. 2008. Are white-tailed deer affecting the ecological integrity of the Smith Island complex? Progress and preliminary findings report submitted to The Bald Head Island Conservancy and Village of Bald Head Island. UNCW Department of Biology and Marine Biology, Wilmington, NC. 44p. Available: <<http://www.bhic.org/wildlife>>. Accessed 9 Jul. 2014.
- CAROLINA VEGETATION SURVEY. 2013. Plot data for Smith Island, North Carolina. University of North Carolina, Chapel Hill, NC. Available: <<http://cvs.bio.unc.edu>>. Accessed 9 Jul. 2014.
- CARSON, W. P., J. A. BANTA, A. A. ROYO, AND C. KIRSCHBAUM. 2005. Plant communities growing on boulders in the Allegheny National Forest: evidence for boulders as refugia from deer and as a bioassay of over-browsing. *Nat. Areas J.*, **25**:10–18.

- COKER, W. C. 1918. A visit to Smith Island (North Carolina). *J. Elisha Mitchell Sci. Soc.*, **34**:150–153.
- CONNER, W. H., W. D. MIXON, AND G. W. WOOD. 2005. Maritime forest habitat dynamics on Bulls Island, Cape Romain National Wildlife Refuge, SC, following Hurricane Hugo. *For. Ecology Manage.*, **212**:127–134.
- COOPER, A. W. AND S. SATTERTHWAITE. 1964. Smith Island and the Cape Fear Peninsula: a comprehensive report on an outstanding natural area. Wildlife Preserves, Inc, Raleigh, NC. 80 p.
- CÔTÉ, S. D., T. P. ROONEY, J. TREMBLAY, C. DUSSAULT, AND D. M. WALLER. 2004. Ecological impacts of deer overabundance. *Ann. Rev. Ecol., Evol. Syst.*, **35**:113–147.
- DEER FACTS. 2009. Guide to deer exclosures. DeerFacts.org. Available <<http://www.scatacook.org/Downloads/DeerExclosures/Deer-Exclosure-Report.pdf>>. Accessed 9 Jul. 2014.
- EPSTEIN, M. B., G. A. FELDHAMER, AND R. L. JOYNER. 1983. Predation on white-tailed deer fawns by bobcats, foxes, and alligators: predator assessment. *Proc. Ann. Conf. Southeast. Fish and Wildlife Agencies*, **37**:244–251.
- , ———, R. J. HAMILTON, AND W. G. MOORE. 1985. Home range and mortality of white-tailed deer fawns in coastal South Carolina. *Proc. Ann. Conf. Southeast. Fish and Wildlife Agencies*, **39**:373–379.
- FORRESTER, J. A., D. J. LEOPOLD, AND H. B. UNDERWOOD. 2006. Isolating the effects of white-tailed deer on the vegetation dynamics of a rare maritime American holly forest. *Am. Midl. Nat.*, **156**:135–150.
- GOODRUM, P. D. AND V. H. REID. 1958. Deer browsing in the longleaf pine belt. *Proc. Soc. Am. For.*, **1958**:139–143.
- HARLOW, R. F. AND R. L. DOWNING. 1970. Deer browsing and hardwood regeneration in the southern Appalachians. *J. For.*, **68**:298–300.
- HORSLEY, S. B., S. L. STOUT, AND D. S. DECALESTA. 2003. White-tailed deer impact on the vegetation dynamics of a northern hardwood forest. *Ecol. Appl.*, **13**:98–118.
- KNIGHT, T. M., J. L. DUNN, L. A. SMITH, J. DAVIS, AND S. KALISZ. 2009. Deer facilitate invasive plant success in Pennsylvania forest understory. *Nat. Areas J.*, **29**:110–116.
- LONG, Z. T., T. H. PENDERGAST, AND W. P. CARSON. 2007. The impact of deer on relationships between tree growth and mortality in an old-growth beech-maple forest. *For. Ecol. Manage.*, **252**:230–38.
- LOPAZANSKI, M. J., J. P. EVANS, AND R. E. SHAW. 1988. An assessment of maritime forest resources on the North Carolina coast. Report prepared for the North Carolina Division of Coastal Management, Raleigh, NC. 108 p.
- McSHEA, W. J., H. B. UNDERWOOD, AND J. H. RAPPOLE (eds.). 1997. The science of overabundance: deer and ecology and population management. Smithsonian Institution Press, Washington, D.C. 402 p.
- MILLER, S. K. 1988. Reproductive biology of white-tailed deer on Cumberland Island, Georgia. Master of Science Thesis, University of Georgia, Athens, GA. 114 p.
- NICHOLAS, C., J. WAMBAUGH, T. LADNER, AND J. M. BENNER. 2007. Deer exclosure guidelines (draft). Pennsylvania Department of Conservation and Natural Resources, Bureau of Forestry. Harrisburg, PA. 70 p. Available: <http://www.apps.dcnr.state.pa.us/forestry/sfrmp/documents/TimberRegen_Deer_Fence_Guidelines_4_7_%202004.pdf>. Accessed 9 Jul. 2014.
- NORTH CAROLINA COASTAL RESERVE. 2014. Bald Head Woods. North Carolina Division of Coastal Management, North Carolina Coastal Reserve Program, Raleigh, NC. Available: <<http://www.nccoastalreserve.net>>. Accessed 9 Jul. 2014.
- OOSTING, H. J. 1954. Vegetation of the maritime strand. *Bot. Rev.*, **20**:226–262.
- OSBOURNE, J. S., A. S. JOHNSON, P. E. HALE, R. L. MARCHINTON, C. V. VANSANT, AND J. M. WENTWORTH. 1992. Population ecology of the Blackbeard Island white-tailed deer. *Bull. 26, Tall Timbers Res., Inc.* 108 p.
- PARNELL, J. F. AND D. A. ADAMS. 1970. Smith Island: a resource capability study, interim report. University of North Carolina at Wilmington, Wilmington, NC. 83 p.
- PEET, R. K., T. R. WENTWORTH, AND P. S. WHITE. 1998. A flexible, multipurpose method for recording vegetation composition and structure. *Castanea*, **63**:262–274.
- R DEVELOPMENT CORE TEAM. 2012. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3- 900051-07-0. Available: <<http://www.R-project.org>>. Accessed 9 Jul. 2014.
- RAY, D. K., E. G. BOLEN, AND W. D. WEBSTER. 2001. Characteristics of a barrier island deer population in the southeastern United States. *J. Elisha Mitchell Sci. Soc.*, **117**:133–122.

- ROONEY, T. P. AND D. M. WALLER. 2003. Direct and indirect effects of white-tailed deer in forest ecosystems. *For. Ecol. Manage.*, **181**:165–176.
- ROWLAND, R. D. 1989. Population genetics of white-tailed deer on Cumberland Island, Georgia. Master of Science Thesis, University of Georgia, Athens, GA. 84 p.
- ROYO, A. A. AND W. P. CARSON. 2006. On the formation of dense understory layers in forests worldwide: consequences and implications for forest dynamics, biodiversity, and succession. *Can. J. For. Res.*, **36**:1345–1362.
- RUSSELL, F. L., D. B. ZIPPIN, AND N. L. FOWLER. 2001. Effects of white-tailed deer (*Odocoileus virginianus*) on plants, plant populations and communities: a review. *Am. Midl. Nat.*, **146**:1–26.
- SCHAFALE, M. P. 2012. Guide to the classification of the natural communities of North Carolina (fourth approximation). North Carolina Natural Heritage Program, Raleigh, N.C. 208 p. Available: <<http://www.ncnhp.org/web/nhp/nhp-publications>>. Accessed 9 Jul. 2014
- SHADOW, R. A. 2011. Plant fact sheet for yaupon, *Ilex vomitoria*. USDA-Natural Resources Conservation Service, East Texas Plant Materials Center, Nacogdoches, TX. Available: <http://plants.usda.gov/factsheet/pdf/fs_ilvo.pdf>. Accessed 9 Jul. 2014.
- SHERILL, B. L., A. G. SNIDER, AND C. S. DePERNO. 2010. White-tailed deer on a barrier island: implications for preserving an ecologically important maritime forest. Proceedings of the annual conference of Southeastern associated fish and wildlife agencies, Maggie Valley, NC. Available: <<http://www4.ncsu.edu/~csdepern/documents/PR42-BHIWTDSEAFWA.pdf>>. Accessed 9 Jul. 2014.
- STRANSKY, J. J. 1969. Deer habitat quality for major forest types in the South, p. 42–45. *In*: L. K. Halls (ed.). White-tailed deer in the southern forest habitat. Proceedings of a symposium held at Nacogdoches, Texas. U.S. Forest Service, Washington, D.C. 130 p.
- STROMAYER, K. A. K. AND R. J. WARREN. 1997. Are overabundant deer herds in the eastern United States creating alternative stable states in forest plant communities? *Wildl. Soc. Bull.*, **25**:227–234.
- TANENTZAP, A. J., D. R. BAZELY, S. KOHB, M. TIMCISKAB, E. G. HAGGITHC, T. J. CARLETON, AND D. A. COOMESA. 2011. Seeing the forest for the deer: do reductions in deer-disturbance lead to forest recovery? *Biol. Cons.*, **144**:376–382.
- TURNER, M. G. AND S. P. BRATTON. 1987. Fire, grazing and the landscape heterogeneity of a Georgia barrier island, p. 85–101. *In*: M. G. Turner (ed.). Landscape Heterogeneity and Disturbance. Springer-Verlag, NY. 239 p.
- U.S. FOREST SERVICE. 2012. Silvics manual, volume 2. *Ilex vomitoria*. U.S. Forest Service, Washington, D.C. Available: <http://www.na.fs.fed.us/pubs/silvics_manual/volume_2/persea/borbonia.htm>. Accessed 9 Jul. 2014.
- WARBINGTON-WELLS, M. R. 2004. Live oak (*Quercus virginiana*) recruitment and seedling survivorship in a southern maritime forest. M.S. thesis, Department of Biology, Georgia Southern University, Statesboro, GA. 92 p.
- WEAKLEY, A. S. 2012. Flora of Southern and Mid-Atlantic States. Working draft of 30 Nov. 2012. University of North Carolina Herbarium, Chapel Hill, NC. 1225 p. Available: <<http://www.herbarium.unc.edu/flora.htm>>. Accessed 9 Jul. 2014.
- WEBSTER, C. R., M. A. JENKINS, AND J. H. ROCK. 2005. Long-term response of spring flora to chronic herbivory and deer exclusion in Great Smoky Mountains National Park, USA. *Biol. Cons.*, **125**:297–307.
- WELLS, B. W. 1932. The natural gardens of North Carolina. The University of North Carolina Press, Chapel Hill, NC. 288 p.
- . 1939. A new forest climax: the salt spray climax of Smith Island, N. C. *Bull. Torrey Bot. Club*, **66**:629–634.