

**A GUIDE TO
SALT MARSH PLANTS
COMMON TO NORTH CAROLINA**

v
-

Written by Elizabeth Jean Wilson,
Hampton Mariners Museum

Illustrated by Carolyn Hoss and Mary Ann Nelson
Edited by Kathy Hart
UNC Sea Grant College Program

\$2.00

This work was partially sponsored by the Office of Sea Grant, NOAA, U.S. Department of Commerce, under Grant No. NA81AA-D-00026, and the North Carolina Department of Administration. The U.S. Government is authorized to produce and distribute reprints for governmental purposes notwithstanding any copyright that may appear hereon.

The Hampton Mariners Museum, Beaufort, N.C., is an extension of the N.C. State Museum of Natural History, a division of the N.C. Department of Agriculture.

UNC Sea Grant College Publication UNC SG 81 04 June 1981

the salt marsh habitat

Salt marshes act as a meeting place between the land and the sea. Marshes, extending hundreds of miles along coastal North Carolina, range throughout the intertidal areas of estuaries, creeks and rivers where the water is salty.

Salt marshes form when land vegetation invades shallow water. Vegetation grows in areas that are exposed during low tides. Silt, clay and organic matter become trapped around plants, which gradually elevates the soil. As the elevation increases, the amount of time the soil is submerged decreases. Gradually other plant species are able to grow on the higher ground.

Special features of the habitat are:

- a constantly changing environment, due to rising and falling tides.
- a tidal cycle that changes water levels, salinity and temperature and exchanges salt water twice every 24 hours.
- marsh plants and animals that are alternately wet and dry.
- salinity that varies from high to low because of varying freshwater flow from rains and streams.
- evaporation that leaves salt residues in the soil and on the plants.
- gentle wind and wave action. Storms and boat wakes are destructive, although marsh grasses reduce some of the damaging effects.
- muddy waters due to silt and organic matter suspended in the slow-flowing water.
- slow-flowing water which often limits oxygen supply in the soil and water. Bacterial activity leaves the mud rich in hydrogen sulfide which gives the marsh its characteristic “rotten egg” odor.

the value of salt marshes

Salt marshes are more than just beautiful grass meadows that reach into intertidal areas along the coast. They are a self-maintaining ecosystem of which the importance extends beyond the boundaries of the marsh.

The amount of energy or food stored in salt marshes has been compared to wheat fields. They are described as one of the most productive natural ecosystems in the world.

Plants are the primary producers of food in the marsh as they utilize energy from the sun in their growth and reproduction. Decomposition of salt marsh grass by bacteria and fungi supplies rich nutrients to the marsh system. These nutrients may be washed out in the estuary by tidal action where they help support plankton, which forms an important part of the base for the estuarine food web. Estuaries, in turn, produce animals important to human consumption like oysters, clams, shrimp, scallops and numerous fishes.

plant characteristics

The salt marsh environment imposes limits on the kinds of plants that can survive. Few species of plants can adapt to daily variations in salinity, water depth and wind and wave action. Direct sunlight contributes to the stress by causing water loss and leaving harmful residues of salt. Plants able to tolerate high salinities are called halophytes.

Ways plants adapt to conditions in the salt marsh are:

- waxy, leathery or fleshy leaves**—resist salt damage and retains moisture. Yaupon, wax myrtle, bamboo vine, marsh elder and seaside goldenrod are examples.
- inrolled or cylindrical leaves**—help to minimize moisture loss by having less surface area. Leaves of cordgrass are inrolled; leaves of needlerush are cylindrical.
- succulents**—store salt in succulent stem and leaf tissues. Water is retained to withstand high salt content without stress, as in glasswort and sea blite.
- salt-secreting**—have special glands to remove excess salt in species such as spike grass and cordgrass.
- extensive rhizomes**—collect more soil and organic matter through underground stems and roots and anchor plants like cordgrass and needlerush to the soil. This prevents severe erosion by tides and storms.
- large cells**—retain air, during periods of submer-sion, in stems and rhizomes, especially in cordgrass and needlerush.
- rotating leaves**—avoid the hottest rays of the sun to minimize water loss during the day, as in marsh pennywort.

Most salt-tolerant plants do grow well outside the saltwater environment. But in freshwater habitats or up land areas, they cannot compete successfully with other species. Remember that plants grow where they can, not where they want.

zonation

Special adaptations of salt marsh plants determine where they live in the marsh. Vegetation zones are more or less distinctive due to the complex combination of environmental factors. Factors include elevation above mean water level, soil type, salinity, temperature, depth of water, period of time the plants are submerged and exposed and the tidal range of the area. The marsh may slope several feet from the supratidal to the subtidal zone.

The five marsh zones are:

—**supratidal**—the area above the average high tide mark. Only unusually high tides and storms reach this area, although plants are exposed to salt spray. Indicator plants are wax myrtle, yaupon, marsh elder, salt-meadow hay and cottonbush.


—**salt barren**—elevated depressions infrequently inundated by salt water. The sun evaporates the water leaving a film of salt on the soil. This area is sparsely vegetated by plants that have adapted to desert-like conditions. Indicator species include succulents such as glasswort and sea blite. Spike grass, which secretes salt through glands, also grows here.

—**upper intertidal**—areas inundated with salt water during short periods of each high tide. Plants are submerged less than 50 percent of the time. Indicator species include cordgrass, needlerush and sea ox-eye.

—**lower intertidal**—a lower elevation zone where plants are submerged more than 50 percent of the time. Cordgrass is dominant, and is generally taller than in the upper intertidal zone. Nutrients are more available and there is less moisture loss.

—**subtidal**—an area submerged nearly all the time. Plants growing here include eel grass and algae.

ZONATION OF A SALT MARSH



how to use this key


This key to common salt marsh plants is intended for layman's use. A background in botany is not necessary to use the guide. Terms used throughout the key, such as leaf shapes and leaf margins, are illustrated.

To use the key, first decide whether the plant in question is a shrub, vine, herb or grass. A general description of each plant form is given on the following pages: shrub—page 9, vine—page 13, herb—page 17, grass—page 23.

After choosing a plant form, turn the page to the simple line key. Beginning with the main heading, at each level choose one of two descriptions. Continue to work through the choices until a plant is named. When a name is reached, turn to the page with the corresponding drawing and description to determine if the identification seems correct. If it is not correct, try again.

It is possible to have a plant that is not included in the key, since only the most common plants are given. Two or three scientific names are given in some cases where species are very similar. It is beyond the scope and purpose of this key to be concerned with these differences. If you wish to go further, use the *Manual of the Flora of the Carolinas* by Radford, Ahles and Bell.


Leaf Shapes




Linear




Lanceolate




Elliptic




Ovate




Oblong




**Triangular
Ovate**



**Arrowhead
Shaped**



Lobed



Round


Leaf Margins



Entire



Toothed




Scalloped

shrub



- woody
- perennial; does not die back to the ground, but persists through winter
- usually branches from the base with several main stems, not usually from a single trunk.


Leaf Arrangements



Opposite



Alternate



Basal Rosette

marsh elder
Iva frutescens

- Asteraceae: Aster family
- Leaves: opposite, lanceolate to elliptic, dark green, fleshy, hairy, margins toothed
- Flowers: cream-colored heads encased by fleshy leaves
- Fruits: tiny nutlets, Aug.-Oct.
- Supratidal zone



sea ox-eye
Borrchia frutescens

- Asteraceae: Aster family
- Leaves: opposite, elliptic to ovate, light green, covered with grey hairs, margins often spiny
- Flowers: large yellow heads, daisy-like
- Fruits: tiny nutlets, head is spiny, May-Sept.
- Upper intertidal zone



wax myrtle
Myrica cerifera

- Myricaceae: Myrtle family
- Leaves: alternate, lanceolate to elliptic, evergreen with yellow resinous glands, aromatic, margins toothed
- Flowers: tiny catkins, April
- Fruits: berry-like, waxy, Aug.-Oct.
- Supratidal zone
- Wax from berries was used in colonial times in bayberry candles. Branches were placed upon fish carts to repel flies.



Shrubs

Leaves alternate on stem

Leaves opposite on stem

Leaves not aromatic or light green

Leaves aromatic when crushed, yellow-green; yellow glands appear as tiny dots on the leaf surface; fruit a waxy, whitish berry—wax myrtle, *Myrica cerifera* (page 11)

Leaves light green; margins often spiny; flowers in large heads with yellow petals—sea ox-eye, *Borrchia frutescens* (page 11)

Leaves dark green; fleshy with tooth margins; flowers in small cream-colored heads encased by fleshy leaves; appear to lack petals—marsh elder, *Iva frutescens* (page 11)


Leaves elliptic to lanceolate, light green and fleshy; margins have coarse teeth; fruiting heads appear cottony in autumn—cottonbush, *halimifolia* (page 12)

Leaves elliptic to ovate, dark green, leathery and waxy; margins have rounded teeth; fruit a red berry—yaupon, *Ilex vomitoria* (page 12)

**groundsel tree
cottonbush**

Baccharis halimifolia

- Asteraceae: Aster family
- Leaves: alternate, ovate to elliptic, light green, margins toothed
- Flowers: heads in clusters, cream-colored
- Fruits: tiny nutlets attached to white hairs giving a cottony appearance, Sept.-Oct.
- Supratidal zone



yaupon
Ilex vomitoria

- Aquifoliaceae: Holly family
- Leaves: alternate, ovate to elliptic, dark green, waxy, margins with rounded teeth
- Flowers: tiny white, four petals, male and female flowers on separate plants, March-May
- Fruits: red berries, Oct.-Dec.
- Supratidal zone
- Tea was made from the leaves in colonial days. A stronger drink was used by Indians as a purgative during religious ceremonies.

vine




- trails along ground or climbs on other plants by tendrils
- may be woody or herbaceous

bamboo vine, greenbriar

Smilax laurifolia

- Liliaceae: Lily family
- Stems: green, spiny
- Leaves: evergreen, leathery, oblong to lanceolate
- Flowers: tiny green, three petals, in umbels, July-Aug.
- Fruits: berries bluish-black, Sept.-Oct.
- Woody high-climbing vine, dense thickets in supratidal zone
- Young shoots may be eaten in salads or cooked like asparagus



catbriar, greenbriar

Smilax bona-nox

- Liliaceae: Lily family
- Stems: green, spiny
- Leaves: evergreen, leathery, ovate to lanceolate, often with basal lobes, margins often spiny, usually mottled
- Flowers: tiny green, three petals, in umbels, April-May
- Fruits: berries bluish-black, Sept.-Nov.
- Woody low-climbing vine, dense thickets in supratidal zone
- Young shoots may be eaten in salads or cooked like asparagus



Vines

Leaves not arrowhead-shaped; flowers not pink and bell-shaped

Leaves arrowhead shaped; flowers pink, bell-shaped—bindweed, morning glory, *Ipomoea sagittata* (page 16)

Woody vine growing on shrubs at edge of marsh, often forms dense thickets; stems usually spiny; leaves leathery

Low twining herb growing on marsh grasses; stems never spiny; sap is milky; leaves linear—milkweed vine, *Cynanchum palustre* (page 16)


Leaves ovate to lanceolate; often with a pair of basal lobes; margins often spiny—greenbriar, catbriar, *Smilax bona-nox* (page 15)

Leaves oblong to lanceolate; margins entire—bamboo vine, greenbriar, *Smilax laurifolia* (page 15)

bindweed, morning glory

Ipomoea sagittata

- Convolvulaceae: Morning glory family
- Leaves: arrowhead-shaped
- Flowers: pink, bell-shaped
- Fruit: capsule, July-Sept.
- Trails over ground in supratidal zone



milkweed vine

Cynanchum palustre

- Asclepiadaceae: Milkweed family
- Stems: herbaceous with milky sap
- Leaves: opposite, linear
- Flowers: greenish tinged with rose, five petals, in umbels, June-July
- Fruits: pods slender, seeds attached to silky white hairs, July-Oct.
- Upper intertidal to supratidal zone



herb



- herbaceous; lacks a persistent woody stem; dies back in winter
- may be perennial; dies back to roots, rhizomes or bulbs
- may be annual; entire plant dies after one growing season; propagates only by seed

Leaves scale-like, roundish or triangular-ovate

Leaves scale-like or roundish with scalloped margins

Leaves roundish, umbrella-like; flowers small, white; stems are fleshy underground runners—marsh pennywort, *Hydrocotyle bonariensis* or *H. umbellata* (page 22)

Leaves scale-like, tiny; stems thick and succulent, jointed—glasswort, *Salicornia virginica*, *S. europaea* or *S. bigelovii* (page 22)

Leaves triangular-ovate, margins coarsely toothed

Leaves lobed, 6 to 14 cm. long, light green and hairy; flowers large, pink, Hibiscus-like; shrub-like, 0.3 to 1 meter tall—marsh mallow, seashore mallow, *Kosteletskya virginica* (page 21)

Leaves not lobed, less than 5 cm. long, light green tinged with purple; flowers green, inconspicuous; grows less than 0.3 meters tall—seabeach orach, *Atriplex patula* (page 22)

Leaves linear, lanceolate or elliptic

Flowering heads daisy-like—white, lavender or yellow

Flowering heads yellow, numerous; leaves elliptic, greater than 1 cm. wide—seaside goldenrod, *Solidago sempervirens* (page 20)

Flowering heads white to lavender, few to many; leaves linear, less than 1 cm. wide—marsh aster, *Aster tenuifolius* or *A. subulatus* (page 20)

Flowers not daisy-like; lavender, pink or greenish

Flowers lavender or pink; leaves not fleshy


Flowers pink, few, greater than 1 cm. wide; leaves along stem, not in basal rosettes—marsh pink, *Sabatia stellaris* (page 21)

Flowers lavender, numerous, less than 1 cm. wide; leaves in basal rosettes—sea lavender, *Limonium carolinianum* or *L. nashii* (page 21)

Flowers greenish, inconspicuous; leaves fleshy—sea blite, *Suaeda linearis* (page 20)

seaside goldenrod
Solidago sempervirens

- Asteraceae: Aster family
- Perennial
- Leaves: elliptic to lanceolate, fleshy, basal, margins toothed, along stem
- Flowers: heads yellow, daisy-like, arranged in rows
- Fruits: tiny nutlets attached to white hairs, Aug.-Nov.
- Supratidal zone
- A herbal tea is made from the leaves and flowers



marsh aster

Aster tenuifolius, *A. subulatus*

- Asteraceae: Aster family
- Annual or perennial
- Leaves: linear, fleshy
- Flowers: heads daisy-like, white or lavender with pale yellow centers, few to many
- Fruits: tiny nutlets attached to white hairs, Sept.-Nov.
- Upper intertidal zone



sea-blite


Suaeda linearis

- Chenopodiaceae: Goosefoot family
- Leaves: alternate, linear, dull green to whitish
- Flowers: green, inconspicuous
- Fruits: nutlet-like, Aug.-frost
- Upper intertidal zone and salt barrens
- Succulent leaves may be eaten raw in salads

sea lavender

Limonium carolinianum, *L. nashii*

- Plumbaginaceae: Sea lavender family
- Leaves: elliptic, fleshy, basal rosettes
- Flowers: small, lavender, five petals, arranged in fan-shaped inflorescence
- Fruits: nutlet-like, Aug.-Oct.
- Upper intertidal zone



marsh pink

Sabatia stellaris

- Gentianaceae: Gentian family
- Stems: Cylindrical or four-sided
- Annual
- Leaves: opposite, lanceolate to elliptic
- Flowers: large, showy, pink with yellow centers, five petals
- Fruit: capsule, July-Aug.
- Upper intertidal zone



seashore mallow, marsh mallow

Kosteletskya virginica

- Malvaceae: Hibiscus family
- Perennial, shrub-like to 1 meter tall
- Leaves: large, triangular-ovate, often lobed, hairy
- Flowers: large, pink, five petals, stamens very prominent
- Fruits: capsule, July-Aug.
- Supertidal zone
- Original "marsh mallow" was made from the roots. Young leaves may be added to thicken stews.



seabeach orach

Atriplex patula

- Chenopodiaceae: Goosefoot family
- Leaves: triangular-ovate, fleshy, light green tinged with purple, margins toothed; lower leaves opposite, upper leaves alternate
- Flowers: green, inconspicuous
- Fruits: nutlet-like, July-frost
- Supratidal to upper intertidal zones, salt barrens



marsh pennywort

Hydrocotyle bonariensis, *H. umbellata*


- Apiaceae: Carrot family
- Stems: underground white rhizomes
- Leaves: roundish, umbrella-like with petiole attached to center, margins scalloped
- Flowers: small, white, arranged in umbels
- Fruits: nutlet-like, similar to caraway or dill seeds, April-Sept.
- Grows low to ground, often forms mats in supratidal zone
- Leaves change orientation during the day to avoid the hottest rays of the sun



glasswort

Salicornia virginica, *S. europaea*, *S. bigelovii*

- Chenopodiaceae: Goosefoot family
- Stems: thick, succulent, jointed, cactus-like, may turn pink in autumn
- Leaves: small, scale-like, opposite
- Flowers: green, inconspicuous
- Fruits: nutlet-like, July-Oct.
- May form mats in upper intertidal zone or salt barrens
- Succulent stems may be eaten in salads




grass



- grasses fall into three distinct families: true grasses, sedges and rushes
- leaves are usually linear and often rolled inward.
- flowers and fruits are small and parts are inconspicuous



black needlerush
Juncus roemerianus

- Juncaceae: Rush family
- Stems: underground rhizomes
- Leaves: stem-like, cylindrical, stiff, spine-tipped, grey-green
- Flowers: clusters of six-petaled, brown, scale-like flowers appear 15 cm. from tip of leaf
- Fruits: tiny brown capsules, May-Oct.
- Grows in dense stands in upper intertidal zone
- The sharp needles were used as sewing needles during colonial times



salt marsh cord grass
saltwater cord grass
Spartina alterniflora


- Poaceae: True grass family
- Stems: cylindrical, hollow, jointed, extensive rhizomes in mud
- Leaves: linear, flat, 4 to 12 mm. wide, margins with tiny sharp teeth near tip
- Flowers: tiny scale-like florets in spikelets, ascending
- Fruits: tiny grains, rice-like, June-Sept.
- Resembles a field of grain; most important plant in intertidal zone; plants often stunted in upper intertidal zone
- Coastal Indians wove mats and baskets with this plant



salt-meadow hay

Spartina patens

- Poaceae: True grass family
- Stems: cylindrical, hollow, jointed, rhizomes in sand
- Leaves: linear, rolled inward, 1 to 6 mm. wide, margins may have tiny sharp teeth near tip
- Flowers: tiny scale-like florets in spikelets, ascending, purplish
- Fruits: tiny grains, June-Sept.
- Similar to salt marsh cord grass, but more narrow and delicate
- Supratidal to upper intertidal zone
- Harvested for cattle feed during colonial times



salt or spike grass

Distichlis spicata

- Poaceae: True grass family
- Stems: cylindrical, hollow, jointed, rhizomes in sand, less than 0.5 meter tall
- Leaves: linear, flat or rolled inward, along opposite sides of stem, flat in one plane, margins smooth
- Flowers: dense clusters of green spikelets
- Druits: grains, June-Oct.
- Forms mats in supratidal, upper intertidal or salt barren zone

marsh sedge

Limbristylis spadicosa

- Cyperaceae: Sedge family
- Stems: triangular
- Leaves: basal, linear, usually rolled inward
- Flowers: oval spikelets, scale-like, in loose clusters
- Fruits: tiny brown nutlets, two-sided, July-Sept.
- Supratidal to upper intertidal zone



three square

Scirpus robustus

- Cyperaceae: Sedge family
- Stems: triangular, sharply angled
- Leaves: along stem, rolled inward
- Flowers: oval to cylindrical spikelets, scale-like, dense clusters below tip of leaf
- Fruits: brown nutlets, two-sided, July-Sept.
- Supratidal to upper intertidal zone



References

- Radford, A. E., H. E. Ahles and C. R. Bell. 1968. *Manual of the Vascular Flora of the Carolinas*. UNC Press, UNC Chapel Hill, N.C.
- Ross, Theodore. 1981. *Intertidal Salt Marshes of Oregon*. Oregon State University Extension Service. Marine Advisory Program, A Land Grant/Sea Grant Cooperative SG 63.
- Shuster, Carl N. Jr. 1966. *The Nature of a Tidal Marsh*. Information Leaflet, N.Y. State Department of Environment and Conservation. Division of Educational Services.
- Silberhorn, Gene M. 1976. *Tidal Wetland Plants of Virginia*. Educational Series No. 19. Virginia Institute of Marine Sciences.
- Spitsbergen, Judith M. 1980. *Seacoast Life: an ecological guide to natural seashore communities in North Carolina*. N.C. Museum of Natural History and Hampton Mariners Museum.
- Taylor, Beth. 1970. *The Field Approach to Coastal Ecology*. Regional Marine Science Project of Carteret County Public Schools.

Index To Common Names

Bamboo vine	15
Bindweed	16
Black Needlerush	25
Catbriar	15
Cottonbush	12
Glasswort	22
Greenbriar	15
Groundsel Tree	12
Marsh Aster	20
Marsh Elder	11
Marsh Mallow	21
Marsh Pennywort	22
Marsh Pink	21
Marsh Sedge	27
Milkweed Vine	16
Morning Glory	16
Salt Grass	26
Spike Grass	26
Salt Marsh Cord Grass	25
Salt Meadow Hay	26
Saltwater Cord Grass	25
Seabeach Orach	22
Sea Blite	20
Sea Lavender	21
Sea Ox-Eye	11
Seaside Goldenrod	20
Seashore Mallow	21
Three Square	27
Wax Myrtle	11
Yaupon	12

Index To Scientific Names

<i>Aster subulatus</i>	20
<i>Aster tenuifolius</i>	20
<i>Atriplex patula</i>	22
<i>Baccharis halimifolia</i>	12
<i>Borrchia frutescens</i>	11
<i>Cynanchum palustre</i>	16
<i>Distichlis spicata</i>	26
<i>Fimbristylis spadicea</i>	27
<i>Hydrocotyle bonariensis</i>	22
<i>Hydrocotyle umbellata</i>	22
<i>Ilex vomitoria</i>	12
<i>Ipomoea sagittata</i>	16
<i>Iva frutescens</i>	11
<i>Juncus roemerianus</i>	25
<i>Kosteletskya virginica</i>	21
<i>Limonium carolinianum</i>	21
<i>Limonium nashii</i>	21
<i>Myrica cerifera</i>	11
<i>Sabatia stellaris</i>	21
<i>Salicornia bigelovii</i>	22
<i>Salicornia europaea</i>	22
<i>Salicornia virginica</i>	22
<i>Scirpus robustus</i>	27
<i>Smilax bona-nox</i>	15
<i>Smilax laurifolia</i>	15
<i>Solidago sempervirens</i>	20
<i>Spartina alterniflora</i>	25
<i>Spartina patens</i>	26
<i>Suaeda linearis</i>	20

Acknowledgements

The author wishes to thank the following people who aided in the writing and publication of this guide.

To the people who reviewed and tested the key: Judith M. Spitsbergen and JoAnne Powell, curators of education at the Hampton Mariners Museum; Mike Alford, curator of maritime research and naturalist at the Hampton Mariners Museum; Bruce Weaver, chief naturalist at the Cape Lookout National Seashore; Mark Joyner, education coordinator for the N.C. Marine Resources Center at Bogue Banks; Ken Moore, superintendent of the N.C. Botanical Garden; Julie Moore, botanist for the N.C. Natural Heritage Program; and to the teachers and students who took the manuscript into the field.

To Carolyn Hoss for her illustrations and testing of the key in the field and to Mary Ann Nelson for use of illustrations from *Seacoast Life: an ecological guide to natural seashore communities in North Carolina* by Judith M. Spitsbergen.

To the UNC Sea Grant College Program, B.J. Copeland, director, for financial assistance and Lundie Mauldin, marine education specialist, for suggesting a need for a key to salt marsh plants for teachers and students.

To Charles McNeill, curator of the Hampton Mariners Museum, for his support of this project.